

Table 4A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 1
Run Date: 3 / 29 / 04

Comparison of Racial and Ethnic Distribution by Clinic							
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Agape WIC	1,584	1.0	98.7	0.1	0.1	0.1	0.0
Albion Cli	767	65.1	31.3	3.3	0.3	0.1	0.0
Alcona	195	98.5	0.5	0.5	0.0	0.5	0.0
Algonac Cl	461	94.4	2.6	1.7	1.3	0.0	0.0
Allegan	369	93.0	3.3	3.3	0.5	0.0	0.0
Alma WIC	192	91.1	3.1	5.7	0.0	0.0	0.0
Alpena Cou	745	99.2	0.5	0.1	0.1	0.0	0.0
Antrim - B	229	94.3	1.3	2.2	1.7	0.4	0.0
Arab/Chald	720	86.1	12.9	0.3	0.0	0.7	0.0
Arenac Cou	429	96.3	0.9	1.6	0.5	0.7	0.0
Associated	43	*	*	*	*	*	*
Baldwin	400	54.5	27.3	16.5	0.0	1.8	0.0
Bangor	1,794	55.8	10.8	33.1	0.2	0.1	0.0
Baraga Cl	190	98.4	0.0	0.5	1.1	0.0	0.0
Barry-Eato	1,484	95.8	2.2	1.5	0.2	0.3	0.0
Barry-Eato	1,156	95.2	1.0	2.9	0.6	0.4	0.0
Battle Cre	3,231	63.2	25.2	10.5	0.4	0.8	0.0
Baxter Com	120	8.3	79.2	7.5	0.8	4.2	0.0
Bay Cl	2,721	94.9	2.5	2.4	0.0	0.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by Clinic							
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Bayside WI	70	*	*	*	*	*	*
Beaver Isl	6	*	*	*	*	*	*
Belding Cl	317	89.0	1.6	7.9	0.6	0.9	0.0
Bell Memor	333	96.4	1.5	0.3	0.9	0.9	0.0
Bendle/Bur	194	92.3	0.5	5.2	2.1	0.0	0.0
Benton Har	2,238	35.3	62.1	1.3	0.4	1.0	0.0
Benzie Cli	373	93.6	0.8	4.8	0.8	0.0	0.0
Berrien -	1,126	75.9	19.4	3.0	0.6	1.0	0.0
Bldhd Migr	25	*	*	*	*	*	*
Branch-Hil	1,146	92.5	0.8	6.6	0.0	0.1	0.0
Branch-Hil	880	97.0	1.5	1.3	0.0	0.2	0.0
Branch-Hil	1,103	68.1	1.2	30.7	0.0	0.0	0.0
Browning-C	70	*	*	*	*	*	*
Bryant Com	130	61.5	22.3	3.1	0.0	13.1	0.0
Burton	4,447	75.6	18.5	4.8	0.2	0.8	0.0
Calumet WI	383	97.9	1.3	0.5	0.3	0.0	0.0
Capac Clin	141	89.4	0.7	9.2	0.0	0.7	0.0
Caro WIC S	24	*	*	*	*	*	*
Cedar Spri	943	89.1	4.1	5.6	0.7	0.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 3
Run Date: 3 / 29 / 04

Comparison of Racial and Ethnic Distribution by Clinic							
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Charlevoix	661	96.5	1.1	1.5	0.5	0.5	0.0
Chass Heal	3,272	6.1	5.5	88.3	0.0	0.1	0.0
Cheboygan	641	97.5	0.9	0.8	0.0	0.8	0.0
Cherry Str	1,057	8.0	19.9	68.6	0.5	3.0	0.0
Chesaning	106	94.3	0.9	4.7	0.0	0.0	0.0
Clare Coun	822	99.3	0.2	0.5	0.0	0.0	0.0
Clinica Sa	909	0.3	0.2	99.2	0.0	0.2	0.0
Corner Hea	518	40.2	56.2	2.5	0.2	1.0	0.0
Crawford	404	98.0	1.2	0.5	0.0	0.2	0.0
Dewitt WIC	267	90.6	6.4	2.6	0.4	0.0	0.0
Dickinson-	805	98.8	0.9	0.0	0.2	0.1	0.0
Dickinson-	335	97.3	0.6	1.2	0.9	0.0	0.0
DMC Prim	212	0.0	99.5	0.0	0.5	0.0	0.0
DMC Prim	54	*	*	*	*	*	*
DMC Prim	25	*	*	*	*	*	*
Dorr Outre	400	97.5	1.3	1.3	0.0	0.0	0.0
Douglas He	2,785	14.9	55.0	23.6	0.3	6.2	0.0
Dowagiac	1,437	81.6	9.4	7.9	0.4	0.8	0.0
Durand WIC	326	97.5	0.0	2.1	0.3	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 4
Run Date: 3 / 29 / 04

Comparison of Racial and Ethnic Distribution by Clinic							
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
East Area	1,319	1.9	96.8	0.7	0.2	0.5	0.0
Eastside H	1,985	3.1	95.1	0.5	0.4	0.9	0.0
Eau Claire	546	2.0	0.7	97.3	0.0	0.0	0.0
Emmet Coun	981	91.9	1.8	1.2	4.8	0.2	0.0
Escanaba C	1,185	98.4	0.3	0.1	0.9	0.3	0.0
Family Com	1,445	37.1	55.0	3.2	0.1	4.6	0.0
Family Hlt	970	39.9	39.5	18.6	0.6	1.4	0.0
Family Roa	738	3.5	90.8	3.4	0.3	2.0	0.0
Farmington	346	77.2	19.9	1.4	0.0	1.4	0.0
Fife Lake	3	*	*	*	*	*	*
Flint Fam	1,327	34.9	61.0	4.1	0.0	0.1	0.0
Gateway	1,561	1.0	98.3	0.3	0.1	0.3	0.0
Gladwin Co	716	99.0	0.8	0.1	0.0	0.0	0.0
Gogebic Cl	325	97.2	0.6	1.8	0.3	0.0	0.0
Golden	140	2.1	97.9	0.0	0.0	0.0	0.0
Grace Prof	293	0.0	100.0	0.0	0.0	0.0	0.0
Grace Ross	2,575	3.6	95.8	0.3	0.1	0.1	0.0
Grand Have	1,012	90.1	1.5	7.6	0.2	0.6	0.0
Grand Trav	22	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 5
Run Date: 3 / 29 / 04

Comparison of Racial and Ethnic Distribution by Clinic							
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Grant	463	65.2	0.6	33.7	0.4	0.0	0.0
Gratiot	1,975	2.0	92.0	0.5	0.2	5.3	0.0
Greenville	436	95.9	0.9	3.2	0.0	0.0	0.0
Hamtramck	29	*	*	*	*	*	*
Hancock Cl	617	94.8	1.6	0.6	0.0	2.9	0.0
Hannahvil	69	*	*	*	*	*	*
Harper Gra	3,124	1.8	97.6	0.3	0.1	0.2	0.0
Haslett Cl	163	77.3	8.6	9.2	0.0	4.9	0.0
Haslett St	18	*	*	*	*	*	*
Herman Kie	3,667	2.4	96.5	0.4	0.1	0.6	0.0
Holland Cl	3,256	47.0	4.0	43.7	0.0	5.3	0.0
Holly WIC	251	85.3	8.4	6.0	0.0	0.4	0.0
Howard Cit	228	98.2	0.9	0.0	0.4	0.4	0.0
Hudsonv	800	89.6	2.0	7.3	0.1	1.0	0.0
Hurley Cli	92	*	*	*	*	*	*
Huron Coun	965	95.0	0.8	3.5	0.4	0.2	0.0
Icchn-Spar	346	0.3	0.0	99.7	0.0	0.0	0.0
Ingham Cou	4,356	55.6	25.8	15.5	0.2	2.9	0.0
Intervale	419	0.2	99.8	0.0	0.0	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by Clinic							
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Ionia Coun	1,020	83.0	2.2	14.2	0.3	0.3	0.0
Iosco	632	96.4	2.5	0.5	0.3	0.3	0.0
Isabella C	1,245	92.2	3.0	3.2	1.0	0.6	0.0
Ithaca Cli	438	95.4	1.6	2.7	0.0	0.2	0.0
Jackson Co	4,460	72.5	20.5	6.1	0.5	0.3	0.0
Janes St A	1,704	11.2	72.8	15.8	0.1	0.1	0.0
Joy-Greenf	3,107	38.9	57.6	3.2	0.1	0.2	0.0
Kalamazoo	2,775	76.8	19.2	3.1	0.1	0.8	0.0
Kalamazoo	1,507	20.8	65.5	12.7	0.2	0.8	0.0
Kalkaska	523	99.8	0.0	0.2	0.0	0.0	0.0
Kent Count	4,642	47.5	33.5	15.8	0.7	2.4	0.0
Kentwood	2,221	43.9	33.4	12.7	0.4	9.7	0.0
Keweenaw	116	8.6	0.0	0.9	88.8	1.7	0.0
Kingsley	60	*	*	*	*	*	*
Lac Vieux	43	*	*	*	*	*	*
Lake Co	335	74.9	21.2	3.3	0.0	0.6	0.0
Lake Odess	112	84.8	0.9	14.3	0.0	0.0	0.0
Lapeer	1,754	96.8	0.6	2.3	0.2	0.1	0.0
Leelanu Cl	246	57.3	1.2	20.7	20.3	0.4	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 7
Run Date: 3 / 29 / 04

Comparison of Racial and Ethnic Distribution by Clinic							
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Leslie-Sto	99	*	*	*	*	*	*
Lincoln Pa	1,488	68.5	24.1	7.2	0.1	0.1	0.0
Livingston	1,088	95.0	1.9	2.3	0.0	0.7	0.0
Lowell	120	85.8	4.2	8.3	1.7	0.0	0.0
M-72 Migra	44	*	*	*	*	*	*
Mancelona	330	98.8	0.6	0.3	0.3	0.0	0.0
Manchester	13	*	*	*	*	*	*
Manchester	46	*	*	*	*	*	*
Manistee C	540	84.4	2.0	7.0	5.7	0.7	0.0
Manistique	308	93.5	0.6	1.0	4.2	0.6	0.0
Mansitee M	31	*	*	*	*	*	*
Marion Bra	169	98.8	0.0	1.2	0.0	0.0	0.0
Marquette	533	92.9	2.4	1.9	2.6	0.2	0.0
Marshall C	256	97.3	0.4	2.3	0.0	0.0	0.0
Mason Clin	111	94.6	1.8	3.6	0.0	0.0	0.0
Mason Co	768	80.3	1.8	16.3	1.2	0.4	0.0
Mccree Hea	5,631	31.6	65.7	2.1	0.1	0.4	0.0
Mcnamara H	248	93.1	2.0	4.4	0.0	0.4	0.0
Mecosta Co	1,148	90.9	6.3	2.0	0.2	0.7	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 8
Run Date: 3 / 29 / 04

Comparison of Racial and Ethnic Distribution by Clinic							
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Medical Pr	47	*	*	*	*	*	*
Medical Pr	37	*	*	*	*	*	*
Menominee	570	99.3	0.2	0.2	0.2	0.2	0.0
Mercy	168	0.0	99.4	0.6	0.0	0.0	0.0
Michigan S	242	35.1	19.8	21.1	0.4	23.6	0.0
Midland Cl	1,649	95.3	3.0	1.2	0.3	0.2	0.0
Midwest	1,766	22.5	35.6	39.3	0.5	2.1	0.0
Migrant H	427	1.9	0.2	97.9	0.0	0.0	0.0
Missaukee	294	99.7	0.0	0.3	0.0	0.0	0.0
Moms & Bab	1,456	8.3	53.0	38.3	0.2	0.3	0.0
Monroe Cou	2,566	90.0	7.1	2.9	0.0	0.0	0.0
Montcalm	37	*	*	*	*	*	*
Montmorenc	278	99.3	0.0	0.4	0.0	0.4	0.0
Mt Clemens	4,208	74.9	17.2	5.8	0.3	1.9	0.0
Munising C	208	96.6	0.5	0.5	2.4	0.0	0.0
Muskegon C	5,776	58.9	30.4	9.2	0.9	0.6	0.0
New Center	39	*	*	*	*	*	*
New Haven	309	63.8	9.1	26.9	0.3	0.0	0.0
New Hope	13	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by Clinic							
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Newaygo	982	92.7	1.1	5.5	0.5	0.2	0.0
Newberry C	264	97.0	1.9	0.0	0.8	0.4	0.0
North Camp	500	35.4	5.4	7.8	0.0	51.4	0.0
Northeast	2,901	4.1	85.0	0.4	0.1	10.4	0.0
Northern M	173	91.9	4.0	0.6	3.5	0.0	0.0
Northwest	762	1.2	98.6	0.3	0.0	0.0	0.0
Oakland Co	5,801	41.1	31.6	24.3	0.3	2.7	0.0
Oceana Co	732	82.5	0.8	16.3	0.4	0.0	0.0
Ogemaw	665	98.5	0.3	0.8	0.3	0.2	0.0
Olhsa Star	793	77.4	16.8	1.6	0.6	3.5	0.0
Olhsa WIC	239	24.3	54.0	20.9	0.4	0.4	0.0
Omni Care	38	*	*	*	*	*	*
Omnicare	6	*	*	*	*	*	*
Omnicare-M	123	0.8	98.4	0.0	0.8	0.0	0.0
Ontonagon	143	95.8	0.0	2.1	1.4	0.7	0.0
Oscoda	205	96.6	1.5	2.0	0.0	0.0	0.0
Otsego Co	786	99.1	0.3	0.1	0.1	0.4	0.0
Park Medic	87	*	*	*	*	*	*
Park Medic	3	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by Clinic							
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Park Medic	64	*	*	*	*	*	*
Paw Paw Cl	568	85.9	4.6	9.2	0.2	0.2	0.0
PCOC	263	87.1	9.1	2.7	0.0	1.1	0.0
Pinconning	148	100.0	0.0	0.0	0.0	0.0	0.0
Port Huron	3,149	85.2	8.1	5.0	1.2	0.5	0.0
Portland	117	95.7	0.9	2.6	0.9	0.0	0.0
Presque Is	299	98.0	0.3	0.7	0.0	1.0	0.0
Profession	126	0.8	99.2	0.0	0.0	0.0	0.0
Pullman	1,029	53.5	4.3	42.0	0.0	0.2	0.0
Reed City	583	99.0	0.7	0.3	0.0	0.0	0.0
Region II	2,021	67.9	5.8	25.8	0.1	0.4	0.0
Riverview	99	*	*	*	*	*	*
Roosevelt-	274	17.9	59.5	22.3	0.4	0.0	0.0
Roscommon	576	98.6	0.5	0.5	0.2	0.2	0.0
S & S Medi	27	*	*	*	*	*	*
Saginaw Co	4,497	51.3	37.8	10.7	0.1	0.2	0.0
Saginaw-Ch	33	*	*	*	*	*	*
Sanilac Co	1,106	96.6	0.5	2.5	0.1	0.3	0.0
Sault Ste	903	64.7	1.8	1.2	31.8	0.6	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 11
Run Date: 3 / 29 / 04

Comparison of Racial and Ethnic Distribution by Clinic							
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Sawyer Afb	281	92.5	3.6	1.8	2.1	0.0	0.0
Schoolcraf	1,681	0.7	98.2	0.4	0.2	0.4	0.0
Seton Cent	170	0.6	99.4	0.0	0.0	0.0	0.0
Shaftsburg	157	98.1	0.6	1.3	0.0	0.0	0.0
Shelby M	478	1.9	0.0	97.9	0.0	0.2	0.0
Sheldon Co	2,677	5.5	50.5	43.5	0.0	0.6	0.0
Shiawassee	1,381	95.1	1.4	2.7	0.5	0.3	0.0
Southfield	3,504	53.9	42.2	1.8	0.1	2.1	0.0
Sparta Pub	236	83.1	4.2	11.4	0.4	0.8	0.0
St Charles	271	97.4	0.0	2.6	0.0	0.0	0.0
St Ignace	260	59.2	1.9	0.8	37.7	0.4	0.0
St Johns W	521	92.7	3.6	3.3	0.2	0.2	0.0
St Louis C	209	96.7	0.0	3.3	0.0	0.0	0.0
St.Clair S	1,758	84.1	11.9	1.6	0.2	2.1	0.0
Stanton Cl	902	95.8	0.9	3.2	0.1	0.0	0.0
Stockbridg	63	*	*	*	*	*	*
Sumpter He	1,275	73.0	25.4	1.3	0.1	0.2	0.0
Taylor	2,899	78.5	20.2	1.2	0.0	0.1	0.0
Teen Healt	7	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by Clinic							
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Temperance	74	*	*	*	*	*	*
Thompsonv	65	*	*	*	*	*	*
Three Oaks	164	90.2	5.5	3.0	0.0	1.2	0.0
Three Rive	1,050	87.0	11.1	1.5	0.1	0.3	0.0
Traverse C	1,826	89.8	2.2	5.4	1.5	1.1	0.0
Tri-Cnty H	708	97.5	0.8	1.6	0.1	0.0	0.0
Tuscola	1,560	98.8	0.8	0.3	0.0	0.1	0.0
U Of M-Ped	31	*	*	*	*	*	*
Vcms River	155	0.6	99.4	0.0	0.0	0.0	0.0
Vicksburg	142	97.2	0.0	2.8	0.0	0.0	0.0
Walled Lak	1,300	78.7	8.2	10.8	0.4	1.9	0.0
Warren Cli	2,683	77.8	13.2	1.8	0.4	6.7	0.0
Washtenaw	127	57.5	33.9	3.1	0.0	5.5	0.0
Wayne-Dear	3,196	98.1	0.9	0.7	0.1	0.2	0.0
Webbervill	62	*	*	*	*	*	*
West Genes	824	27.7	46.8	19.8	0.1	5.6	0.0
Western Wa	2,090	72.8	24.2	2.6	0.1	0.3	0.0
Westland C	2,454	64.8	31.8	2.5	0.2	0.7	0.0
Westside C	2,098	31.0	10.6	56.6	0.7	1.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 4A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Racial and Ethnic Distribution by Clinic							
Clinic	Number Included in Analysis (3)	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaskan Native %	Asian/Pacific Islander %	All Other / Unknown %
Westwood	133	74.4	18.8	6.0	0.0	0.8	0.0
Wexford	1,093	98.6	0.4	0.5	0.1	0.5	0.0
Wheeler WI	25	*	*	*	*	*	*
Whitmore L	61	*	*	*	*	*	*
Willow	16	*	*	*	*	*	*
Woodside	263	58.6	1.9	1.5	37.6	0.4	0.0
Wyoming CI	2,176	61.9	14.3	19.9	0.6	3.4	0.0
Yale Clini	168	97.0	1.2	1.8	0.0	0.0	0.0
Ypsilanti	3,432	42.8	47.2	7.5	0.1	2.4	0.0
Michigan	222,370	57.7	29.1	11.1	0.7	1.4	0.0
Nation (Prior Year)	5,333,226	39.1	20.8	32.2	1.2	3.3	3.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with errors.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 1
Run Date: 3 / 29 / 04

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Agape WIC	1,584	40.2	18.1	41.7
Albion Cli	767	36.9	20.3	42.8
Alcona	195	35.4	22.6	42.1
Algonac Cl	461	39.5	17.8	42.7
Allegan	369	39.3	18.7	42.0
Alma WIC	192	36.5	13.5	50.0
Alpena Cou	745	30.9	20.4	48.7
Antrim - B	229	36.7	17.5	45.9
Arab/Chald	720	30.8	20.3	48.9
Arenac Cou	429	34.5	25.4	40.1
Associated	43	*	*	*
Baldwin	400	46.0	19.3	34.8
Bangor	1,794	29.2	22.6	48.2
Baraga Cl	190	30.5	17.9	51.6
Barry-Eato	1,484	37.9	18.9	43.3
Barry-Eato	1,156	33.0	20.4	46.6
Battle Cre	3,231	40.8	19.7	39.5
Baxter Com	120	24.2	22.5	53.3

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 2
Run Date: 3 / 29 / 04

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Bay Cl	2,721	34.0	21.7	44.3
Bayside WI	70	*	*	*
Beaver Isl	6	*	*	*
Belding Cl	317	37.5	18.0	44.5
Bell Memor	333	34.2	19.5	46.2
Bendle/Bur	194	27.8	17.5	54.6
Benton Har	2,238	29.9	22.6	47.5
Benzie Cli	373	38.3	17.7	44.0
Berrien -	1,126	34.4	21.0	44.6
Bldhd Migr	25	*	*	*
Branch-Hil	1,146	37.8	18.4	43.8
Branch-Hil	880	26.4	24.0	49.7
Branch-Hil	1,103	37.7	19.0	43.2
Browning-C	70	*	*	*
Bryant Com	130	29.2	20.8	50.0
Burton	4,447	41.1	19.6	39.3
Calumet WI	383	31.1	16.7	52.2
Capac Clin	141	40.4	14.2	45.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 3
Run Date: 3 / 29 / 04

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Caro WIC S	24	*	*	*
Cedar Spri	943	33.5	21.2	45.3
Charlevoix	661	33.4	20.4	46.1
Chass Heal	3,272	32.4	25.9	41.8
Cheboygan	641	27.5	22.5	50.1
Cherry Str	1,057	33.0	21.9	45.0
Chesaning	106	27.4	19.8	52.8
Clare Coun	822	32.2	23.4	44.4
Clinica Sa	909	44.2	17.5	38.3
Corner Hea	518	48.1	23.2	28.8
Crawford	404	36.4	19.3	44.3
Dewitt WIC	267	36.3	18.7	44.9
Dickinson-	805	31.3	20.1	48.6
Dickinson-	335	35.5	19.7	44.8
DMC Prim	212	23.1	18.9	58.0
DMC Prim	54	*	*	*
DMC Prim	25	*	*	*
Dorr Outre	400	35.3	21.3	43.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 4
Run Date: 3 / 29 / 04

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Douglas He	2,785	34.7	21.4	43.9
Dowagiac	1,437	35.7	18.8	45.5
Durand WIC	326	34.7	21.8	43.6
East Area	1,319	36.0	19.1	44.9
Eastside H	1,985	32.6	21.3	46.0
Eau Claire	546	41.8	17.8	40.5
Emmet Coun	981	35.9	18.5	45.7
Esanaba C	1,185	28.9	21.2	50.0
Family Com	1,445	37.9	20.0	42.1
Family Hlt	970	41.0	17.1	41.9
Family Roa	738	43.8	21.5	34.7
Farmington	346	37.6	13.9	48.6
Fife Lake	3	*	*	*
Flint Fam	1,327	41.1	17.0	41.8
Gateway	1,561	34.5	18.6	46.8
Gladwin Co	716	31.4	19.1	49.4
Gogebic Cl	325	40.0	17.2	42.8
Golden	140	20.7	27.1	52.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 5
Run Date: 3 / 29 / 04

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Grace Prof	293	24.2	27.3	48.5
Grace Ross	2,575	41.6	21.1	37.3
Grand Have	1,012	34.2	20.1	45.8
Grand Trav	22	*	*	*
Grant	463	31.7	23.8	44.5
Gratiot	1,975	35.2	22.3	42.4
Greenville	436	38.8	19.5	41.7
Hamtramck	29	*	*	*
Hancock Cl	617	35.7	18.3	46.0
Hannahvil	69	*	*	*
Harper Gra	3,124	37.9	24.5	37.5
Haslett Cl	163	32.5	22.7	44.8
Haslett St	18	*	*	*
Herman Kie	3,667	43.5	18.8	37.7
Holland Cl	3,256	32.9	21.6	45.5
Holly WIC	251	35.1	17.5	47.4
Howard Cit	228	34.2	19.3	46.5
Hudsonv	800	33.5	22.6	43.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 6
Run Date: 3 / 29 / 04

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Hurley Cli	92	*	*	*
Huron Coun	965	34.0	18.7	47.4
Iocchn-Spar	346	28.3	24.9	46.8
Ingham Cou	4,356	33.8	22.6	43.6
Intervale	419	22.4	22.0	55.6
Ionia Coun	1,020	38.2	20.1	41.7
Iosco	632	34.0	19.5	46.5
Isabella C	1,245	38.4	20.8	40.8
Ithaca Cli	438	43.6	16.4	40.0
Jackson Co	4,460	37.0	20.6	42.4
Janes St A	1,704	29.6	19.0	51.4
Joy-Greenf	3,107	32.2	21.0	46.9
Kalamazoo	2,775	38.9	17.9	43.2
Kalamazoo	1,507	34.0	18.5	47.5
Kalkaska	523	39.2	19.7	41.1
Kent Count	4,642	31.9	21.6	46.5
Kentwood	2,221	36.9	21.0	42.1
Keweenaw	116	24.1	20.7	55.2

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 7
Run Date: 3 / 29 / 04

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Kingsley	60	*	*	*
Lac Vieux	43	*	*	*
Lake Co	335	33.4	20.9	45.7
Lake Odess	112	38.4	19.6	42.0
Lapeer	1,754	37.5	21.6	40.9
Leelanau Cl	246	32.1	24.8	43.1
Leslie-Sto	99	*	*	*
Lincoln Pa	1,488	42.1	22.6	35.3
Livingston	1,088	40.2	20.6	39.2
Lowell	120	34.2	22.5	43.3
M-72 Migra	44	*	*	*
Mancelona	330	28.8	22.1	49.1
Manchester	13	*	*	*
Manchester	46	*	*	*
Manistee C	540	35.7	19.1	45.2
Manistique	308	26.9	22.1	51.0
Mansitee M	31	*	*	*
Marion Bra	169	34.3	19.5	46.2

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 8
Run Date: 3 / 29 / 04

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Marquette	533	34.9	20.6	44.5
Marshall C	256	42.2	20.7	37.1
Mason Clin	111	39.6	18.0	42.3
Mason Co	768	30.9	21.2	47.9
Mccree Hea	5,631	39.4	18.3	42.3
Mcnamara H	248	37.5	19.0	43.5
Mecosta Co	1,148	33.4	22.5	44.2
Medical Pr	47	*	*	*
Medical Pr	37	*	*	*
Menominee	570	30.9	22.8	46.3
Mercy	168	23.2	23.2	53.6
Michigan S	242	28.1	28.1	43.8
Midland Cl	1,649	33.8	21.8	44.3
Midwest	1,766	30.7	19.8	49.5
Migrant H	427	29.0	18.7	52.2
Missaukee	294	29.3	22.4	48.3
Moms & Bab	1,456	35.2	24.6	40.2
Monroe Cou	2,566	38.6	21.2	40.2

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 9
Run Date: 3 / 29 / 04

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Montcalm	37	*	*	*
Montmorenc	278	29.1	20.5	50.4
Mt Clemens	4,208	42.9	21.5	35.6
Munising C	208	26.9	24.0	49.0
Muskegon C	5,776	28.9	23.0	48.1
New Center	39	*	*	*
New Haven	309	46.3	22.0	31.7
New Hope	13	*	*	*
Newaygo	982	34.5	20.0	45.5
Newberry C	264	30.3	19.3	50.4
North Camp	500	37.8	22.2	40.0
Northeast	2,901	38.9	21.0	40.1
Northern M	173	36.4	17.9	45.7
Northwest	762	34.8	18.6	46.6
Oakland Co	5,801	36.6	21.1	42.3
Oceana Co	732	34.7	19.1	46.2
Ogemaw	665	32.3	20.6	47.1
Olhsa Star	793	43.3	18.2	38.6

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 10
Run Date: 3 / 29 / 04

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Olhsa WIC	239	38.1	22.6	39.3
Omni Care	38	*	*	*
Omnicare	6	*	*	*
Omnicare-M	123	26.0	24.4	49.6
Ontonagon	143	35.7	20.3	44.1
Oscoda	205	38.0	19.5	42.4
Otsego Co	786	24.0	24.0	51.9
Park Medic	87	*	*	*
Park Medic	3	*	*	*
Park Medic	64	*	*	*
Paw Paw Cl	568	32.4	22.2	45.4
PCOC	263	32.7	22.4	44.9
Pinconning	148	37.8	18.9	43.2
Port Huron	3,149	36.1	19.9	44.0
Portland	117	49.6	17.1	33.3
Presque Is	299	30.8	25.1	44.1
Profession	126	11.1	19.8	69.0
Pullman	1,029	30.7	19.4	49.9

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 11
Run Date: 3 / 29 / 04

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Reed City	583	38.3	20.1	41.7
Region II	2,021	39.9	19.7	40.4
Riverview	99	*	*	*
Roosevelt-	274	27.7	15.3	56.9
Roscommon	576	28.8	25.2	46.0
S & S Medi	27	*	*	*
Saginaw Co	4,497	33.2	22.4	44.5
Saginaw-Ch	33	*	*	*
Sanilac Co	1,106	36.5	20.3	43.1
Sault Ste	903	35.4	18.5	46.1
Sawyer Afb	281	34.2	21.4	44.5
Schoolcraf	1,681	38.3	19.0	42.7
Seton Cent	170	27.1	29.4	43.5
Shaftsborg	157	29.9	21.7	48.4
Shelby M	478	25.9	20.7	53.3
Sheldon Co	2,677	28.8	19.1	52.2
Shiawassee	1,381	30.8	23.0	46.3
Southfield	3,504	39.5	20.7	39.8

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 12
Run Date: 3 / 29 / 04

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Sparta Pub	236	36.9	17.8	45.3
St Charles	271	35.1	17.3	47.6
St Ignace	260	30.4	20.8	48.8
St Johns W	521	36.5	21.9	41.7
St Louis C	209	46.9	13.9	39.2
St.Clair S	1,758	44.8	21.7	33.5
Stanton Cl	902	36.8	18.1	45.1
Stockbridg	63	*	*	*
Sumpter He	1,275	42.0	21.1	36.9
Taylor	2,899	41.8	21.8	36.4
Teen Healt	7	*	*	*
Temperance	74	*	*	*
Thompsonv	65	*	*	*
Three Oaks	164	40.9	18.3	40.9
Three Rive	1,050	41.3	18.8	39.9
Traverse C	1,826	34.7	21.1	44.2
Tri-Cnty H	708	36.9	20.1	43.1
Tuscola	1,560	36.9	20.6	42.5

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 13
Run Date: 3 / 29 / 04

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
U Of M-Ped	31	*	*	*
Vcms River	155	23.2	19.4	57.4
Vicksburg	142	31.7	15.5	52.8
Walled Lak	1,300	44.9	19.2	35.9
Warren Cli	2,683	43.0	20.8	36.2
Washtenaw	127	36.2	22.0	41.7
Wayne-Dear	3,196	34.8	22.1	43.1
Webbervill	62	*	*	*
West Genes	824	27.4	19.5	53.0
Western Wa	2,090	41.6	18.9	39.5
Westland C	2,454	47.7	20.9	31.5
Westside C	2,098	30.9	20.6	48.5
Westwood	133	26.3	33.1	40.6
Wexford	1,093	41.4	16.6	42.1
Wheeler WI	25	*	*	*
Whitmore L	61	*	*	*
Willow	16	*	*	*
Woodside	263	31.2	19.8	49.0

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 5A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Age Distribution by Clinic
Children Aged < 5 Years ⁽²⁾

Page: 14
Run Date: 3 / 29 / 04

Comparison of Age Distribution by Clinic				
Clinic	Number Included in Analysis (3,4)	0 - 11 Months %	12 - 23 Months %	24 - 59 Months %
Wyoming Cl	2,176	37.2	20.5	42.3
Yale Clini	168	36.3	20.2	43.5
Ypsilanti	3,432	38.4	19.9	41.7
Michigan	222,370	36.0	20.7	43.2
Nation (Prior Year)	5,333,226	36.6	21.7	41.7

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Records rejected if date of birth and date of visit unknown.

* Percentages are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 1
Run Date: 3 / 29 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Agape WIC	404	8.9 (58)	7.7 (40)	1,529	7.8(14	2.9(62)	14.7(15	639	13.9(29)	12.4(71)		845	18.5 (144)
Albion Cli	189	9.5 (77)	5.3 (16)	747	8.2(15	5.0(13	10.0(31)	318	15.7(54)	10.7(39)		458	9.4 (72)
Alcona	24	* *	* *	191	6.3(91)	8.9(18	5.2 (2)	85	* *	* *		117	11.1 (90)
Algonac Cl	134	9.0 (64)	10.4 (95)	449	5.1(38)	4.5(12	11.1(57)	192	18.8(12	13.5(86)		281	5.0 (25)
Allegan	80	* *	* *	353	7.9(14	6.8(17	13.9(13	148	12.8(19)	8.8(20)		229	9.2 (70)
Alma WIC	43	* *	* *	190	8.4(15	2.6(43)	24.2(19	95	* *	* *		91	* *
Alpena Cou	162	7.4 (33)	9.3 (76)	731	5.1(38)	1.6(11)	13.5(12	354	16.1(63)	14.7(11		493	8.3 (58)
Antrim - B	48	* *	* *	221	7.7(14	4.1(10	14.5(15	107	15.9(57)	15.9(13		140	2.9 (10)
Arab/Chald	143	6.3 (20)	8.4 (54)	706	4.8(31)	3.7(90)	14.3(14	339	18.6(11	14.7(11		460	17.8 (142)
Arenac Cou	95	* *	* *	419	6.2(89)	3.1(71)	13.6(12	169	13.6(23)	11.8(57)		284	9.9 (76)
Associated	6	* *	* *	43	* *	* *	* *	18	* *	* *		30	* *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 2
Run Date: 3 / 29 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Baldwin	125	10.4 (90)	6.4 (24)	391	10.5(18	3.8(96)	9.5(21)	141	11.3(14)	12.1(62)		205	13.7 (116)
Bangor	235	8.5 (51)	8.1 (45)	1,746	8.1(15	4.5(12	15.5(15)	848	17.7(10)	16.3(14)		1,192	12.3 (105)
Baraga Cl	42	* *	* *	184	5.4(48)	3.3(77)	9.8(27)	95	* *	* *		105	2.9 (10)
Barry-Eato	332	9.0 (64)	9.0 (70)	1,444	10.3(18	2.7(49)	12.9(10)	623	22.6(15)	14.3(10)		892	4.1 (20)
Barry-Eato	238	8.8 (56)	7.1 (34)	1,121	5.4(48)	5.6(15	11.5(72)	528	19.1(12)	11.9(59)		749	7.3 (46)
Battle Cre	877	9.5 (77)	7.9 (44)	3,129	9.8(18	3.1(71)	14.4(14)	1,254	18.3(11)	15.6(13)		1,906	18.9 (147)
Baxter Com	13	* *	* *	118	2.5 (3)	5.1(14	13.6(12)	61	* *	* *		96	* *
Bay Cl	595	7.6 (35)	10.9 (102)	2,650	9.0(16	2.3(30)	12.5(89)	1,177	15.5(51)	14.0(97)		1,674	10.7 (85)
Bayside WI	6	* *	* *	67	* *	* *	* *	45	* *	* *		58	* *
Beaver Isl	1	* *	* *	6	* *	* *	* *	3	* *	* *		1	* *
Belding Cl	66	* *	* *	312	5.4(48)	2.2(25)	10.6(49)	135	16.3(67)	14.1(10)		213	8.5 (61)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
Bell Memor	77	* *	* *	325	9.2(17)	2.2(25)	13.8(13)	148	25.0(15)	16.2(13)	194 7.7 (52)
Bendle/Bur	38	* *	* *	189	6.9(11)	4.2(11)	11.1(57)	100	19.0(12)	11.0(43)	128 24.2 (159)
Benton Har	388	12.6 (105)	7.0 (31)	2,207	6.7(10)	3.0(68)	12.5(89)	1,052	14.7(40)	11.4(52)	1,565 25.0 (162)
Benzie Cli	97	* *	* *	365	7.7(14)	6.3(17)	15.6(16)	162	13.6(23)	17.3(14)	221 7.7 (52)
Berrien -	247	7.7 (38)	10.5 (96)	1,105	4.3(24)	1.4 (9)	10.8(51)	486	14.2(33)	10.1(34)	712 13.8 (117)
Bldhd Migr	3	* *	* *	24	* *	* *	* *	13	* *	* *	14 * *
Branch-Hil	287	8.0 (40)	11.1 (105)	1,127	7.8(14)	4.7(12)	13.0(10)	490	16.3(67)	14.5(11)	646 9.1 (68)
Branch-Hil	108	9.3 (73)	4.6 (6)	870	5.4(48)	5.2(14)	10.9(52)	433	18.5(11)	13.6(89)	663 8.4 (60)
Branch-Hil	218	9.2 (72)	5.5 (19)	1,075	9.4(17)	2.7(49)	24.5(19)	462	20.1(13)	35.1(15)	683 18.2 (143)
Browning-C	17	* *	* *	70	* *	* *	* *	29	* *	* *	43 * *
Bryant Com	14	* *	* *	125	4.0(18)	10.4(19)	12.8(97)	64	* *	* *	95 * *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 4
Run Date: 3 / 29 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
Burton	1,030	9.0 (64)	8.6 (60)	4,391	4.8(31)	4.9(13)	11.3(64)	1,737	17.4(94)	14.0(97)	2,649 21.8 (151)
Calumet WI	70	* *	* *	375	7.5(13)	4.3(11)	14.9(15)	192	23.4(15)	16.7(14)	208 2.4 (5)
Capac Clin	38	* *	* *	132	6.8(10)	6.1(16)	16.7(17)	59	* *	* *	86 * *
Caro WIC S	3	* *	* *	22	* *	* *	* *	12	* *	* *	19 * *
Cedar Spri	237	6.8 (25)	11.0 (104)	932	6.0(79)	3.4(80)	8.9(15)	417	17.3(93)	11.3(51)	590 1.9 (1)
Charlevoix	144	7.6 (35)	4.9 (8)	643	3.4(13)	5.0(13)	15.7(16)	296	19.9(13)	14.9(12)	445 3.6 (16)
Chass Heal	788	7.2 (30)	10.0 (91)	3,175	9.2(17)	4.8(13)	13.3(11)	1,311	12.0(17)	12.4(71)	2,089 13.1 (113)
Cheboygan	102	2.9 (1)	13.7 (117)	631	5.9(71)	3.5(81)	12.7(93)	323	14.9(43)	15.8(13)	456 10.5 (83)
Cherry Str	244	6.6 (23)	8.2 (48)	1,040	3.2(12)	5.2(14)	10.9(52)	474	14.3(37)	15.4(12)	706 7.2 (44)
Chesaning	12	* *	* *	102	8.8(16)	2.9(62)	15.7(16)	55	* *	* *	76 * *
Clare Coun	167	10.8 (94)	6.6 (26)	812	5.9(71)	4.8(13)	11.3(64)	355	16.6(78)	14.4(10)	528 11.6 (96)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 5
Run Date: 3 / 29 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Clinica Sa	277	5.1 (11)	6.9 (30)	879	11.4(19)	3.3(77)	20.4(18)	338	18.6(11)	25.7(15)		529	8.5 (61)
Corner Hea	179	9.5 (77)	8.4 (54)	510	4.1(20)	7.5(18)	10.4(40)	148	14.2(33)	12.2(65)		275	14.9 (129)
Crawford	102	3.9 (7)	10.8(100)	395	5.3(46)	2.3(30)	11.4(70)	179	16.8(82)	15.6(13)		234	8.5 (61)
Dewitt WIC	51	* *	* *	258	8.9(16)	4.7(12)	13.2(11)	118	15.3(49)	14.4(10)		160	8.1 (56)
Dickinson-	167	7.2 (30)	13.2 (116)	781	6.9(11)	1.7(13)	12.7(93)	372	17.2(90)	14.0(97)		541	8.5 (61)
Dickinson-	74	* *	* *	326	4.6(29)	4.0(10)	11.3(64)	154	20.8(14)	15.6(13)		214	6.5 (38)
DMC Prim	24	* *	* *	197	5.1(38)	4.1(10)	15.2(15)	114	16.7(79)	9.6(26)		161	32.3 (171)
DMC Prim	6	* *	* *	53	* *	* *	* *	27	* *	* *		42	* *
DMC Prim	5	* *	* *	25	* *	* *	* *	6	* *	* *		19	* *
Dorr Outre	89	* *	* *	392	11.2(18)	4.1(10)	16.8(17)	170	18.2(11)	15.3(12)		260	10.4 (81)
Douglas He	616	8.4 (48)	8.9 (69)	2,710	6.3(91)	7.0(17)	12.8(97)	1,185	8.1 (7)	6.2 (9)		1,595	12.8(111)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 6
Run Date: 3 / 29 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
Dowagiac	330	9.1 (71)	7.3 (36)	1,375	6.6(10)	6.5(17)	11.6(73)	631	14.4(38)	10.9(42)	871 13.1 (113)
Durand WIC	82	* *	* *	308	7.5(13)	4.2(11)	10.1(34)	140	20.7(14)	11.4(52)	209 5.3 (29)
East Area	213	8.9 (58)	5.2 (14)	1,270	9.4(17)	5.9(16)	16.7(17)	584	18.0(10)	12.3(66)	522 31.2 (169)
Eastside H	409	16.4 (117)	4.4 (4)	1,924	9.1(17)	8.5(18)	13.1(11)	872	6.9 (4)	5.5 (4)	1,319 23.6 (156)
Eau Claire	143	3.5 (4)	10.5 (96)	516	13.0(19)	4.7(12)	20.9(18)	207	16.9(84)	24.6(15)	283 15.5 (133)
Emmet Coun	207	5.3 (12)	5.3 (16)	947	3.8(16)	3.7(90)	13.2(11)	431	15.8(55)	13.5(86)	635 7.7 (52)
Escanaba C	245	8.2 (41)	12.2 (110)	1,150	5.5(53)	2.7(49)	10.3(36)	575	19.5(13)	14.1(10)	765 2.1 (3)
Family Com	275	8.4 (48)	7.3 (36)	1,390	6.8(10)	7.3(17)	12.2(83)	592	10.8(11)	14.4(10)	697 29.4 (165)
Family Hlt	243	13.2 (111)	7.8 (42)	944	6.0(79)	7.1(17)	10.3(36)	397	15.9(57)	9.6(26)	555 24.5 (160)
Family Roa	224	17.0 (118)	7.6 (38)	711	2.8 (8)	8.2(18)	8.6(13)	252	13.1(21)	7.9(14)	442 36.4 (173)
Farmington	90	* *	* *	339	7.1(12)	5.6(15)	9.7(25)	163	16.0(61)	8.6(19)	219 11.0 (89)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 7
Run Date: 3 / 29 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
Fife Lake	0	* *	* *	3	* *	* *	* *	1	* *	* *	3 * *
Flint Fam	326	9.5 (77)	5.5 (19)	1,307	6.5(95)	3.9(10)	12.9(10)	542	16.4(72)	14.9(12)	796 27.5 (164)
Gateway	275	15.6 (116)	4.7 (7)	1,500	10.0(18)	5.3(14)	15.6(16)	716	16.8(82)	16.3(14)	757 19.7 (148)
Gladwin Co	135	4.4 (8)	9.6 (81)	702	7.7(14)	1.0 (3)	13.2(11)	348	18.1(10)	12.4(71)	471 3.4 (15)
Gogebic Cl	87	* *	* *	321	7.5(13)	3.1(71)	8.1(11)	141	14.2(33)	7.1(11)	178 10.7 (85)
Golden	13	* *	* *	134	6.0(79)	5.2(14)	16.4(16)	68	* *	* *	116 37.1 (174)
Grace Prof	25	* *	* *	285	2.5 (3)	6.3(17)	10.5(46)	141	10.6(10)	5.0 (3)	231 29.4 (165)
Grace Ross	635	12.0 (99)	6.0 (21)	2,517	7.0(12)	4.1(10)	12.0(78)	943	20.1(13)	13.0(82)	1,534 11.3 (92)
Grand Have	224	10.7 (92)	8.5 (56)	980	5.8(66)	4.4(11)	12.8(97)	463	17.7(10)	11.2(46)	625 8.5 (61)
Grand Trav	1	* *	* *	22	* *	* *	* *	16	* *	* *	18 * *
Grant	102	5.9 (16)	10.8 (100)	450	11.3(19)	2.7(49)	16.0(16)	205	18.5(11)	11.2(46)	285 17.2 (138)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 8
Run Date: 3 / 29 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Gratiot	460	14.8 (113)	7.0 (31)	1,936	7.9(14	4.0(10	12.0(78)	818	14.1 (32)	9.7(29)		1,067	12.7 (110)
Greenville	111	6.3 (20)	9.0 (70)	430	4.2(22)	4.7(12	10.0(31)	183	14.8(42)	7.7(12)		270	6.7 (40)
Hamtramck	0	* *	* *	28	* *	* *	* *	21	* *	* *		21	* *
Hancock Cl	148	6.1 (18)	12.2 (110)	603	4.5(26)	3.8(96)	7.6 (5)	280	7.5 (5)	4.3 (2)		336	3.0 (12)
Hannahvil	11	* *	* *	68	* *	* *	* *	33	* *	* *		44	* *
Harper Gra	830	15.3 (115)	6.3 (22)	2,986	10.0(18	10.8(19	9.6(23)	1,123	6.4 (2)	4.1 (1)		1,455	14.4 (126)
Haslett Cl	23	* *	* *	155	5.2(44)	2.6(43)	14.2(14	70	* *	* *		112	8.9 (67)
Haslett St	2	* *	* *	18	* *	* *	* *	3	* *	* *		11	* *
Herman Kie	1,035	14.9 (114)	4.2 (3)	3,514	8.6(16	7.8(18	10.5(46)	1,338	6.2 (1)	5.8 (6)		2,071	21.2 (149)
Holland Cl	555	10.5 (91)	8.8 (68)	3,147	9.0(16	2.3(30)	14.3(14	1,445	17.4(94)	14.8(11		2,048	5.5 (31)
Holly WIC	52	* *	* *	245	6.9(11	4.5(12	10.2(35)	114	16.7(79)	12.3(66)		155	10.3 (79)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 9
Run Date: 3 / 29 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Howard Cit	47	* *	* *	228	6.6(10	2.2(25)	15.4(15	109	17.4(94)	13.8(91)	156	14.1 (122)
Hudsonv	138	12.3 (101)	8.7 (62)	760	8.7(16	5.9(16	10.0(31)	343	16.0(61)	8.2(16)	513	5.8 (33)
Hurley Cli	2	* *	* *	88	* *	* *	* *	35	* *	* *	62	* *
Huron Coun	202	8.9 (58)	12.4 (113)	949	5.2(44)	4.4(11	11.3(64)	443	14.0(30)	11.1 (44)	607	7.6 (49)
Icchn-Spar	66	* *	* *	317	9.5(18	3.8(96)	18.6(18	157	15.9(57)	21.0(15	238	7.6 (49)
Ingham Cou	747	10.0 (86)	9.0 (70)	4,247	6.8(10	3.6(84)	12.7(93)	1,852	17.5(98)	14.0(97)	2,970	12.5 (108)
Intervale	50	* *	* *	411	8.8(16	2.7(49)	22.4(19	226	23.9(15	23.0(15	322	44.7 (176)
Ionia Coun	257	7.0 (27)	9.3 (76)	996	5.0(36)	5.2(14	10.6(49)	414	20.0(13	13.5(86)	635	6.6 (39)
Iosco	80	* *	* *	622	4.8(31)	3.7(90)	14.0(13	289	20.4(14	14.5(11	407	11.3 (92)
Isabella C	312	8.3 (45)	9.3 (76)	1,224	6.5(95)	2.9(62)	11.4(70)	504	16.5(74)	14.3(10	795	6.8 (42)
Ithaca Cli	112	12.5 (103)	7.1 (34)	425	8.2(15	1.9(16)	20.7(18	174	22.4(15	28.2(15	216	4.6 (24)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Jackson Co	1,165	8.2 (41)	8.7 (62)	4,324	7.4(13)	5.6(15)	12.8(97)	1,822	18.7(12)	14.5(11)		2,437	5.3 (29)
Janes St A	353	12.5 (103)	3.7 (2)	1,648	7.4(13)	2.6(43)	16.7(17)	871	16.2(65)	12.9(79)		1,185	17.6 (141)
Joy-Greenf	630	9.7 (84)	8.7 (62)	3,033	5.9(71)	3.7(90)	11.3(64)	1,444	17.1 (88)	12.0(60)		1,989	31.4 (170)
Kalamazoo	692	9.4 (75)	9.5 (80)	2,679	8.5(15)	4.7(12)	14.0(13)	1,176	17.7(10)	15.4(12)		1,567	22.8 (152)
Kalamazoo	332	13.0 (110)	7.8 (42)	1,464	5.6(57)	6.6(17)	13.7(12)	695	14.7 (40)	15.1(12)		1,015	30.9 (168)
Kalkaska	137	6.6 (23)	5.1 (12)	514	6.8(10)	2.9(62)	10.9(52)	213	18.8(12)	12.7 (77)		338	13.9 (119)
Kent Count	883	9.9 (85)	9.7 (87)	4,487	7.5(13)	2.7(49)	13.0(10)	2,080	16.3(67)	12.9(79)		2,960	1.9 (1)
Kentwood	538	8.7 (55)	10.0 (91)	2,183	6.0(79)	2.9(62)	12.9(10)	919	18.8(12)	13.3(83)		1,268	3.2 (13)
Keweenaw	25	* *	* *	115	4.3(24)	4.3(11)	13.9(13)	64	* *	* *		74	* *
Kingsley	8	* *	* *	60	* *	* *	* *	37	* *	* *		46	* *
Lac Vieux	11	* *	* *	43	* *	* *	* *	22	* *	* *		24	* *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 11
Run Date: 3 / 29 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Lake Co	82	* *	* *	328	7.0(12	5.5(15	11.9(77)	150	24.7(15	10.7(39)	209	13.9 (119)
Lake Odess	30	* *	* *	112	8.0(14	3.6(84)	14.3(14	46	* *	* *	71	* *
Lapeer	442	7.2 (30)	11.1 (105)	1,717	9.4(17	3.3(77)	13.9(13	706	18.3(11	16.3(14	1,087	9.5 (73)
Leelanu Cl	43	* *	* *	243	8.6(16	1.6(11)	19.8(18	105	17.1 (88)	19.0(15	167	12.0 (100)
Leslie-Sto	9	* *	* *	96	* *	* *	* *	45	* *	* *	73	* *
Lincoln Pa	410	8.3 (45)	9.0 (70)	1,455	5.8(66)	2.1(20)	12.2(83)	514	19.5(13	12.1(62)	856	4.1 (20)
Livingston	290	10.3 (89)	16.2 (118)	1,069	6.1(87)	3.6(84)	12.1(82)	416	11.1 (13)	8.9(22)	665	2.7 (9)
Lowell	13	* *	* *	119	5.9(71)	0.8 (1)	13.4(12	54	* *	* *	69	* *
M-72 Migra	8	* *	* *	43	* *	* *	* *	19	* *	* *	29	* *
Mancelona	59	* *	* *	319	5.6(57)	5.0(13	13.5(12	156	20.5(14	16.0(13	222	5.0 (25)
Manchester	2	* *	* *	12	* *	* *	* *	6	* *	* *	7	* *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
Manchester	4	* *	* *	43	* *	* *	* *	22	* *	* *	34 * *
Manistee C	140	13.6 (112)	8.6 (60)	520	6.5 (95)	2.7 (49)	10.4 (40)	240	17.5 (98)	13.3 (83)	334 7.5 (47)
Manistique	62	* *	* *	300	2.7 (6)	9.7 (19)	9.7 (25)	149	20.1 (13)	13.4 (85)	190 17.4 (140)
Mansitee M	6	* *	* *	31	* *	* *	* *	20	* *	* *	24 * *
Marion Bra	21	* *	* *	165	6.7 (10)	4.2 (11)	10.9 (52)	77	* *	* *	125 12.0 (100)
Marquette	126	3.2 (3)	11.9 (109)	520	7.1 (12)	1.2 (5)	11.2 (59)	232	20.3 (14)	13.8 (91)	304 10.9 (88)
Marshall C	68	* *	* *	249	6.4 (94)	4.4 (11)	16.5 (17)	92	* *	* *	121 8.3 (58)
Mason Clin	30	* *	* *	109	5.5 (53)	2.8 (59)	17.4 (17)	46	* *	* *	67 * *
Mason Co	182	4.9 (10)	9.9 (89)	748	3.1 (11)	5.5 (15)	7.6 (5)	356	9.0 (8)	9.0 (23)	492 12.4 (106)
Mccree Hea	1,432	12.2 (100)	5.2 (14)	5,484	5.7 (61)	4.3 (11)	10.4 (40)	2,321	16.4 (72)	11.5 (54)	3,318 25.7 (163)
Mcnamara H	59	* *	* *	239	7.1 (12)	2.1 (20)	13.4 (12)	106	15.1 (46)	16.0 (13)	163 6.7 (40)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
Mecosta Co	256	5.9 (16)	12.5 (114)	1,131	4.2(22)	5.6(15)	8.0 (8)	502	15.3(49)	8.8(20)	745 17.3 (139)
Medical Pr	4	* *	* *	47	* *	* *	* *	21	* *	* *	34 * *
Medical Pr	2	* *	* *	35	* *	* *	* *	24	* *	* *	26 * *
Menominee	118	7.6 (35)	8.5 (56)	563	4.1(20)	3.0(68)	8.0 (8)	259	17.4(94)	8.5(17)	363 3.9 (17)
Mercy	15	* *	* *	161	5.6(57)	3.1(71)	22.4(19)	82	* *	* *	127 38.6 (175)
Michigan S	33	* *	* *	236	2.1 (1)	5.5(15)	4.2 (1)	106	17.0(87)	8.5(17)	173 9.2 (70)
Midland Cl	366	7.1 (28)	9.6 (81)	1,614	8.6(16)	2.3(30)	17.7(18)	719	16.3(67)	11.8(57)	1,084 10.6 (84)
Midwest	331	12.7 (106)	7.6 (38)	1,674	7.6(14)	6.3(17)	17.1(17)	841	18.1(10)	16.6(14)	295 9.8 (75)
Migrant H	87	* *	* *	412	5.1(38)	1.2 (5)	23.1(19)	214	18.2(11)	23.4(15)	272 5.1 (27)
Missaukee	60	* *	* *	288	4.5(26)	2.4(37)	7.6 (5)	145	17.2(90)	9.7(29)	192 13.5 (115)
Moms & Bab	384	10.2 (88)	6.8 (29)	1,386	13.9(19)	4.8(13)	21.1(18)	569	22.5(15)	23.2(15)	422 6.4 (37)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Monroe Cou	675	8.4 (48)	8.3 (51)	2,506	7.3(12	2.0(18)	12.8(97)	1,006	18.1(10	13.8(91)		1,408	14.7 (128)
Montcalm	8	* *	* *	35	* *	* *	* *	21	* *	* *		29	* *
Montmorenc	58	* *	* *	274	5.5(53)	5.8(15)	10.9(52)	138	8.0 (6)	13.8(91)		187	7.5 (47)
Mt Clemens	1,167	9.0 (64)	8.7 (62)	4,131	5.7(61)	2.9(62)	14.0(13	1,472	13.7(25)	9.5(25)		2,378	11.9 (98)
Munising C	34	* *	* *	202	4.5(26)	1.0 (3)	17.3(17	97	* *	* *		154	9.1 (68)
Muskegon C	1,192	9.0 (64)	8.1 (45)	5,581	6.0(79)	5.3(14)	10.3(36)	2,736	16.5(74)	12.0(60)		3,387	15.3 (131)
New Center	4	* *	* *	38	* *	* *	* *	20	* *	* *		33	* *
New Haven	84	* *	* *	301	8.3(15	1.7(13)	20.6(18	99	* *	* *		168	11.9 (98)
New Hope	1	* *	* *	13	* *	* *	* *	6	* *	* *		9	* *
Newaygo	232	8.6 (54)	10.3 (94)	941	7.3(12	3.6(84)	11.2(59)	422	13.3(22)	9.7(29)		603	22.9 (153)
Newberry C	54	* *	* *	255	5.1(38)	3.5(81)	17.6(17	128	18.0(10	18.8(14		171	11.1 (90)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
North Camp	115	7.8 (39)	9.6 (81)	491	3.9(17)	5.5(15)	5.9 (4)	200	11.0(12)	5.5 (4)	280 17.1 (137)
Northeast	645	12.9 (109)	5.0 (10)	2,787	8.6(16)	9.1(18)	14.2(14)	1,104	6.8 (3)	5.9 (7)	1,593 13.9 (119)
Northern M	41	* *	* *	173	7.5(13)	1.2 (5)	13.3(11)	78	* *	* *	98 * *
Northwest	122	12.3 (101)	8.2 (48)	744	6.6(10)	5.9(16)	9.4(20)	347	13.8(28)	7.8(13)	236 18.6 (145)
Oakland Co	1,398	9.6 (82)	8.3 (51)	5,699	6.1(87)	3.1(71)	11.2(59)	2,395	18.6(11)	15.5(12)	3,689 14.3 (125)
Oceana Co	166	9.0 (64)	6.6 (26)	710	8.5(15)	1.3 (8)	12.7(93)	321	20.2(13)	11.2(46)	471 10.0 (77)
Ogemaw	131	3.8 (6)	9.9 (89)	653	5.7(61)	4.0(10)	12.3(86)	311	15.8(55)	11.6(55)	420 12.9 (112)
Olhse Star	237	11.0 (95)	5.1 (12)	775	5.9(71)	4.6(12)	8.9(15)	305	11.5(16)	9.2(24)	447 10.3 (79)
Olhse WIC	53	* *	* *	230	10.4(18)	4.8(13)	10.4(40)	97	* *	* *	145 24.1 (158)
Omni Care	0	* *	* *	38	* *	* *	* *	23	* *	* *	31 * *
Omnicare	0	* *	* *	6	* *	* *	* *	4	* *	* *	4 * *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
Omnicare-M	11	* *	* *	113	2.7 (6)	6.2(16)	8.0 (8)	63	* *	* *	92 * *
Ontonagon	35	* *	* *	142	3.5(14)	3.5(81)	8.5(12)	66	* *	* *	89 * *
Oscoda	48	* *	* *	203	6.9(11)	4.9(13)	11.3(64)	84	* *	* *	120 14.2 (123)
Otsego Co	109	12.8 (108)	10.1 (93)	765	2.9(10)	6.1(16)	9.9(28)	404	12.1 (18)	10.6(38)	568 2.6 (7)
Park Medic	3	* *	* *	87	* *	* *	* *	53	* *	* *	71 * *
Park Medic	0	* *	* *	3	* *	* *	* *	3	* *	* *	1 * *
Park Medic	6	* *	* *	62	* *	* *	* *	32	* *	* *	50 * *
Paw Paw Cl	74	* *	* *	549	8.6(16)	6.2(16)	11.7(75)	256	18.8(12)	11.7(56)	353 21.5 (150)
PCOC	53	* *	* *	255	5.9(71)	4.3(11)	12.5(89)	115	19.1(12)	14.8(11)	166 15.1 (130)
Pinconning	38	* *	* *	144	8.3(15)	2.1(20)	10.4(40)	61	* *	* *	74 * *
Port Huron	755	9.3 (73)	5.0 (10)	3,056	5.8(66)	7.9(18)	10.3(36)	1,369	14.9(43)	11.2(46)	1,977 10.2 (78)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
Portland	36	* *	* *	115	7.0(12)	6.1(16)	13.0(10)	41	* *	* *	72 * *
Presque Is	65	* *	* *	293	5.1(38)	2.0(18)	9.2(19)	128	15.6(52)	10.2(35)	206 15.5 (133)
Profession	7	* *	* *	124	4.0(18)	2.4(37)	13.7(12)	87	* *	* *	102 30.4 (167)
Pullman	171	2.9 (1)	6.4 (24)	1,004	7.4(13)	4.3(11)	13.9(13)	496	16.1 (63)	12.9(79)	685 16.4 (136)
Reed City	147	7.5 (34)	12.9 (115)	574	2.8 (8)	3.7(90)	8.7(14)	234	13.7(25)	11.1 (44)	369 12.2 (102)
Region II	523	7.1 (28)	4.4 (4)	1,964	7.2(12)	6.1(16)	14.0(13)	801	14.5(39)	15.6(13)	1,066 15.5 (133)
Riverview	19	* *	* *	98	* *	* *	* *	37	* *	* *	67 * *
Roosevelt-	58	* *	* *	253	3.6(15)	2.8(59)	17.0(17)	151	21.2(14)	9.9(32)	187 24.6 (161)
Roscommon	127	6.3 (20)	8.7 (62)	570	5.8(66)	2.6(43)	11.2(59)	262	13.7(25)	10.7(39)	389 3.3 (14)
S & S Medi	1	* *	* *	27	* *	* *	* *	21	* *	* *	22 * *
Saginaw Co	1,021	9.6 (82)	7.7 (40)	4,260	8.0(14)	2.8(59)	15.4(15)	1,940	18.7(12)	15.1(12)	3,077 6.9 (43)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 18
Run Date: 3 / 29 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
Saginaw-Ch	8	* *	* *	32	* *	* *	* *	12	* *	* *	19 * *
Sanilac Co	261	8.8 (56)	5.4 (18)	1,094	5.0(36)	5.4(14)	9.1(17)	473	15.2(48)	12.7(77)	693 4.2 (23)
Sault Ste	196	3.6 (5)	12.2(110)	893	5.3(46)	2.5(41)	14.7(15)	415	18.3(11)	14.7(11)	607 5.6 (32)
Sawyer Afb	65	* *	* *	276	7.2(12)	1.4 (9)	14.1(14)	124	21.0(14)	12.1(62)	169 2.4 (5)
Schoolcraf	430	10.7 (92)	3.3 (1)	1,644	6.2(89)	2.7(49)	12.2(83)	709	14.0(30)	10.2(35)	1,028 14.4(126)
Seton Cent	5	* *	* *	166	6.0(79)	11.4(19)	12.0(78)	75	* *	* *	140 22.9(153)
Shafterburg	32	* *	* *	147	10.2(18)	8.2(18)	15.6(16)	75	* *	* *	112 10.7 (85)
Shelby M	81	* *	* *	445	6.5(95)	2.7(49)	16.2(16)	238	19.3(13)	22.7(15)	323 12.4(106)
Sheldon Co	501	9.4 (75)	7.0 (31)	2,596	6.7(10)	2.1(20)	15.9(16)	1,339	20.7(14)	17.5(14)	1,857 3.9 (17)
Shiawassee	333	6.9 (26)	9.3 (76)	1,318	5.9(71)	7.4(18)	13.1(11)	635	19.4(13)	15.6(13)	907 6.0 (36)
Southfield	850	11.2 (96)	9.1 (74)	3,445	5.6(57)	5.2(14)	9.1(17)	1,367	9.4 (9)	6.8(10)	2,202 15.3(131)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Sparta Pub	34	* *	* *	233	7.3(12	2.6(43)	11.6(73)	106	14.2(33)	17.9(14		153	2.6 (7)
St Charles	64	* *	* *	260	9.2(17	3.1(71)	16.2(16	125	24.0(15	15.2(12		185	5.9 (35)
St Ignace	53	* *	* *	253	4.7(30)	3.6(84)	13.8(13	128	17.2(90)	17.2(14		173	5.8 (33)
St Johns W	106	5.7 (13)	8.5 (56)	514	2.5 (3)	3.9(10	9.9(28)	212	11.3(14)	6.1 (8)		318	9.7 (74)
St Louis C	53	* *	* *	201	10.0(18	2.5(41)	17.9(18	80	* *	* *		79	* *
St.Clair S	495	9.5 (77)	8.7 (62)	1,730	6.0(79)	2.4(37)	13.3(11	590	15.1(46)	10.2(35)		981	12.2 (102)
Stanton Cl	223	5.8 (14)	4.9 (8)	891	6.6(10	2.4(37)	12.3(86)	393	15.0(45)	9.9(32)		569	10.4 (81)
Stockbridg	6	* *	* *	61	* *	* *	* *	29	* *	* *		39	* *
Sumpter He	335	11.3 (98)	9.6 (81)	1,235	7.5(13	2.3(30)	13.1(11	461	17.8(10	13.9(96)		709	7.6 (49)
Taylor	801	8.9 (58)	9.6 (81)	2,824	9.1(17	4.0(10	10.5(46)	1,034	16.2(65)	13.6(89)		1,622	8.6 (66)
Teen Healt	1	* *	* *	7	* *	* *	* *	4	* *	* *		5	* *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
Temperance	5	* *	* *	72	* *	* *	* *	41	* *	* *	53 * *
Thompsonv	8	* *	* *	63	* *	* *	* *	36	* *	* *	43 * *
Three Oaks	44	* *	* *	158	5.7(61)	1.9(16)	9.5(21)	60	* *	* *	100 8.0 (55)
Three Rive	296	4.4 (8)	8.1 (45)	1,035	7.9(14)	2.3(30)	14.9(15)	418	20.8(14)	12.4(71)	618 5.2 (28)
Traverse C	445	8.3 (45)	11.2 (107)	1,796	6.5(95)	5.1(14)	13.0(10)	805	16.9(84)	12.5(75)	1,110 13.8 (117)
Tri-Cnty H	173	5.8 (14)	11.6 (108)	696	6.3(91)	6.5(17)	12.5(89)	297	21.5(14)	12.5(75)	442 12.2 (102)
Tuscola	394	8.9 (58)	9.6 (81)	1,531	6.9(11)	3.7(90)	12.3(86)	641	15.6(52)	14.2(10)	981 4.1 (20)
U Of M-Ped	6	* *	* *	30	* *	* *	* *	16	* *	* *	12 * *
Vcms River	16	* *	* *	135	2.2 (2)	27.4(19)	5.2 (2)	75	* *	* *	122 33.6 (172)
Vicksburg	27	* *	* *	138	9.4(17)	3.6(84)	13.8(13)	72	* *	* *	101 18.8 (146)
Walled Lak	377	8.5 (51)	8.2 (48)	1,277	6.0(79)	2.7(49)	10.4(40)	467	16.5(74)	9.6(26)	770 11.4 (94)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 21
Run Date: 3 / 29 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Warren Cli	740	8.9 (58)	9.1 (74)	2,636	6.5(95)	2.2(25)	14.8(15)	959	16.5(74)	12.3(66)	1,558	11.4 (94)
Washtenaw	24	* *	* *	121	5.8(66)	7.4(18)	9.9(28)	51	* *	* *	74	* *
Wayne-Dear	695	8.2 (41)	6.3 (22)	3,135	5.4(48)	3.8(96)	11.8(76)	1,352	16.7(79)	14.5(11)	1,957	14.2 (123)
Webbervill	10	* *	* *	62	* *	* *	* *	34	* *	* *	43	* *
West Genes	102	12.7 (106)	9.8 (88)	811	5.7(61)	4.9(13)	12.8(97)	418	16.3(67)	14.1(10)	596	11.6 (96)
Western Wa	576	8.5 (51)	8.3 (51)	1,992	6.5(95)	2.1(20)	14.2(14)	818	19.2(12)	15.2(12)	1,261	24.0 (157)
Westland C	753	11.2 (96)	8.5 (56)	2,399	5.9(71)	2.6(43)	11.2(59)	756	18.0(10)	13.8(91)	1,283	8.2 (57)
Westside C	418	6.2 (19)	10.5 (96)	2,070	5.5(53)	1.7(13)	13.6(12)	987	21.6(15)	14.7(11)	1,421	2.3 (4)
Westwood	22	* *	* *	128	14.8(19)	0.8 (1)	19.5(18)	53	* *	* *	92	* *
Wexford	294	8.2 (41)	10.5 (96)	1,080	4.8(31)	2.2(25)	9.6(23)	454	15.9(57)	12.3(66)	656	12.5 (108)
Wheeler WI	11	* *	* *	24	* *	* *	* *	6	* *	* *	6	* *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
Whitmore L	14	* *	* *	60	* *	* *	* *	21	* *	* *	29 * *
Willow	1	* *	* *	16	* *	* *	* *	5	* *	* *	10 * *
Woodside	57	* *	* *	258	8.5(15)	2.3(30)	12.8(97)	127	22.0(15)	7.9(14)	180 7.2 (44)
Wyoming Cl	542	9.0 (64)	10.9(102)	2,101	6.9(11)	3.0(68)	12.8(97)	892	12.9(20)	11.2(46)	1,217 3.9 (17)
Yale Clini	46	* *	* *	164	4.9(35)	5.5(15)	14.0(13)	72	* *	* *	98 * *
Ypsilanti	790	10.1 (87)	6.6 (26)	3,345	6.8(10)	5.8(15)	12.0(78)	1,411	16.9(84)	12.3(66)	2,113 23.2 (155)
Michigan	50,792	9.4	8.3	216,362	6.9	4.3	12.7	93,962	16.1	12.9	134,250 13.1
Nation (Prior Year)	1,208,400	8.9	7.6	4,913,316	6.3	5.4	13.5	2,069,605	15.4	14.3	3,295,974 13.1

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 1
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Agape WIC	413	43.8 (84)	329	14.0 (52)	206	12.1 (41)
Albion Cli	195	39.0(101)	151	7.3(102)	121	7.4 (84)
Alcona	50	* *	33	* *	32	* *
Algonac Cl	135	48.1 (72)	79	* *	62	* *
Allegan	91	* *	81	* *	55	* *
Alma WIC	48	* *	32	* *	14	* *
Alpena Cou	163	61.3 (22)	129	17.1 (34)	124	13.7 (23)
Antrim - B	54	* *	43	* *	30	* *
Arab/Chald	155	58.1 (31)	103	22.3 (15)	97	* *
Arenac Cou	97	* *	93	* *	96	* *
Associated	6	* *	12	* *	9	* *
Baldwin	134	51.5 (57)	71	* *	48	* *
Bangor	365	53.7 (48)	288	12.8 (59)	298	12.4 (36)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 2
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Baraga Cl	43	* *	35	* *	28	* *
Barry-Eato	368	58.2 (30)	282	11.0 (76)	231	11.3 (48)
Barry-Eato	255	44.3 (82)	233	12.4 (61)	195	10.3 (56)
Battle Cre	906	39.7 (97)	712	11.1 (73)	533	6.2 (92)
Baxter Com	13	* *	19	* *	19	* *
Bay Cl	613	49.6 (66)	460	10.9 (77)	414	7.5 (80)
Bayside WI	5	* *	8	* *	9	* *
Beaver Isl	1	* *	0	* *	0	* *
Belding Cl	66	* *	69	* *	49	* *
Bell Memor	77	* *	58	* *	49	* *
Bendle/Bur	38	* *	26	* *	23	* *
Benton Har	391	37.3(107)	633	10.1 (84)	432	8.8 (66)
Benzie Cli	99	* *	73	* *	47	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 3
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Berrien -	258	45.0 (81)	246	14.2 (51)	210	13.8 (22)
Bldhd Migr	6	* *	4	* *	4	* *
Branch-Hil	307	48.2 (71)	219	15.5 (44)	163	12.3 (38)
Branch-Hil	111	57.7 (33)	165	7.9(100)	166	10.8 (49)
Branch-Hil	281	58.7 (26)	192	20.8 (19)	151	12.6 (34)
Browning-C	18	* *	10	* *	9	* *
Bryant Com	14	* *	24	* *	20	* *
Burton	1,215	38.5(103)	861	10.3 (81)	635	8.3 (71)
Calumet WI	77	* *	52	* *	36	* *
Capac Clin	39	* *	38	* *	21	* *
Caro WIC S	3	* *	4	* *	1	* *
Cedar Spri	239	58.6 (28)	177	15.8 (42)	157	8.3 (71)
Charlevoix	151	61.6 (19)	106	18.9 (23)	100	13.0 (28)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 4
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Chass Heal	821	52.9 (50)	315	17.1 (34)	558	15.6 (19)
Cheboygan	102	67.6 (8)	135	19.3 (22)	106	17.0 (12)
Cherry Str	247	66.8 (9)	190	21.1 (17)	197	16.2 (17)
Chesaning	12	* *	17	* *	14	* *
Clare Coun	181	43.6 (86)	169	8.9 (94)	157	6.4 (90)
Clinica Sa	278	94.2 (1)	200	22.5 (14)	125	20.8 (4)
Corner Hea	181	51.9 (56)	124	8.1 (99)	94	* *
Crawford	106	53.8 (46)	74	* *	58	* *
Dewitt WIC	67	* *	54	* *	34	* *
Dickinson-	169	56.2 (41)	152	16.4 (38)	114	12.3 (38)
Dickinson-	74	* *	53	* *	50	* *
DMC Prim	24	* *	46	* *	26	* *
DMC Prim	6	* *	9	* *	13	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 5
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
DMC Prim	5	* *	7	* *	9	* *
Dorr Outre	93	* *	72	* *	61	* *
Douglas He	633	46.4 (76)	524	22.7 (11)	403	18.9 (5)
Dowagiac	348	52.0 (54)	202	14.9 (46)	186	12.4 (36)
Durand WIC	83	* *	50	* *	44	* *
East Area	321	21.5(119)	202	6.9(106)	179	10.6 (52)
Eastside H	416	35.3(109)	386	8.3 (97)	319	6.3 (91)
Eau Claire	160	76.3 (3)	114	41.2 (3)	77	* *
Emmet Coun	240	66.3 (10)	187	18.2 (26)	141	17.7 (9)
Escanaba C	245	49.4 (70)	205	11.2 (72)	192	10.4 (55)
Family Com	340	41.5 (91)	298	16.8 (36)	238	12.6 (34)
Family Hlt	263	49.8 (65)	181	16.0 (41)	146	8.2 (73)
Family Roa	224	21.9(118)	164	15.2 (45)	109	11.9 (42)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 6
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Farmington	94	* *	50	* *	35	* *
Fife Lake	0	* *	1	* *	2	* *
Flint Fam	351	31.1(114)	305	9.2 (91)	176	8.5 (68)
Gateway	400	21.3(120)	228	7.0(105)	179	3.9 (98)
Gladwin Co	149	59.7 (23)	129	14.0 (52)	117	7.7 (78)
Gogebic Cl	88	* *	64	* *	34	* *
Golden	13	* *	32	* *	31	* *
Grace Prof	25	* *	65	* *	74	* *
Grace Ross	649	40.2 (94)	638	9.9 (87)	378	6.6 (88)
Grand Have	239	54.8 (43)	185	18.9 (23)	162	16.7 (16)
Grand Trav	1	* *	3	* *	4	* *
Grant	106	54.7 (44)	68	* *	81	* *
Gratiot	473	31.5(113)	263	7.2(104)	274	5.1 (96)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 7
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Greenville	112	52.7 (52)	76	* *	51	* *
Hamtramck	0	* *	0	* *	0	* *
Hancock Cl	152	70.4 (6)	104	32.7 (4)	71	* *
Hannahvil	11	* *	15	* *	14	* *
Harper Gra	935	27.0(116)	376	11.4 (70)	517	9.5 (62)
Haslett Cl	26	* *	44	* *	31	* *
Haslett St	2	* *	7	* *	5	* *
Herman Kie	1,119	31.8(112)	759	9.0 (93)	522	7.3 (85)
Holland Cl	734	62.9 (17)	619	22.6 (12)	532	14.7 (20)
Holly WIC	57	* *	29	* *	35	* *
Howard Cit	47	* *	44	* *	24	* *
Hudsonv	188	66.0 (11)	134	20.9 (18)	134	18.7 (6)
Hurley Cli	3	* *	20	* *	11	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 8
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Huron Coun	205	45.4 (80)	185	6.5(108)	132	4.5 (97)
Icchn-Spar	77	* *	66	* *	59	* *
Ingham Cou	823	61.6 (19)	963	15.7 (43)	757	13.2 (24)
Intervale	50	* *	94	* *	78	* *
Ionia Coun	258	50.8 (59)	174	17.2 (33)	166	11.4 (47)
Iosco	128	37.5(106)	124	8.9 (94)	92	* *
Isabella C	316	64.6 (15)	258	9.3 (90)	211	11.8 (44)
Ithaca Cli	129	50.4 (63)	99	* *	46	* *
Jackson Co	1,186	50.0 (64)	1,013	10.9 (77)	696	8.0 (74)
Janes St A	355	33.0(111)	251	6.8(107)	228	7.9 (76)
Joy-Greenf	642	57.5 (35)	542	27.3 (8)	491	32.0 (2)
Kalamazoo	734	40.6 (93)	524	23.5 (10)	339	17.7 (9)
Kalamazoo	351	48.1 (72)	240	8.3 (97)	201	6.5 (89)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 9
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Kalkaska	137	68.6 (7)	101	18.8 (25)	84	* *
Kent Count	916	58.6 (28)	898	16.4 (38)	746	11.8 (44)
Kentwood	550	57.8 (32)	463	16.4 (38)	349	10.6 (52)
Keweenaw	25	* *	14	* *	14	* *
Kingsley	8	* *	8	* *	6	* *
Lac Vieux	11	* *	6	* *	3	* *
Lake Co	86	* *	54	* *	45	* *
Lake Odess	30	* *	18	* *	19	* *
Lapeer	447	51.2 (58)	346	14.7 (48)	293	9.6 (61)
Leelanu Cl	47	* *	47	* *	45	* *
Leslie-Sto	10	* *	20	* *	17	* *
Lincoln Pa	425	43.3 (87)	306	11.1 (73)	251	12.7 (33)
Livingston	297	65.0 (14)	206	20.4 (20)	156	12.8 (31)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 10
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Lowell	14	* *	28	* *	17	* *
M-72 Migra	10	* *	8	* *	5	* *
Mancelona	61	* *	59	* *	67	* *
Manchester	2	* *	2	* *	2	* *
Manchester	4	* *	6	* *	11	* *
Manistee C	142	59.2 (25)	91	* *	87	* *
Manistique	62	* *	49	* *	53	* *
Mansitee M	6	* *	2	* *	2	* *
Marion Bra	37	* *	34	* *	29	* *
Marquette	130	65.4 (13)	93	* *	79	* *
Marshall C	75	* *	52	* *	29	* *
Mason Clin	33	* *	29	* *	14	* *
Mason Co	182	53.8 (46)	121	13.2 (56)	126	10.3 (56)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 11
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Mccree Hea	1,497	31.1(114)	954	6.0(109)	743	3.5 (99)
Mcnamara H	59	* *	38	* *	28	* *
Mecosta Co	264	55.3 (42)	200	12.5 (60)	191	9.9 (59)
Medical Pr	4	* *	13	* *	9	* *
Medical Pr	2	* *	2	* *	8	* *
Menominee	121	39.7 (97)	90	* *	92	* *
Mercy	18	* *	36	* *	28	* *
Michigan S	37	* *	49	* *	50	* *
Midland Cl	368	66.0 (11)	315	16.5 (37)	270	11.9 (42)
Midwest	403	35.7(108)	224	13.8 (55)	225	7.6 (79)
Migrant H	88	* *	58	* *	52	* *
Missaukee	61	* *	42	* *	48	* *
Moms & Bab	417	38.8(102)	224	17.9 (27)	222	18.5 (7)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 12
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Monroe Cou	695	46.0 (78)	568	11.4 (70)	438	7.5 (80)
Montcalm	8	* *	2	* *	5	* *
Montmorenc	60	* *	45	* *	43	* *
Mt Clemens	1,208	47.4 (74)	966	12.3 (63)	683	8.5 (68)
Munising C	34	* *	40	* *	44	* *
Muskegon C	1,216	42.4 (89)	822	11.6 (67)	988	7.5 (80)
New Center	4	* *	7	* *	5	* *
New Haven	88	* *	83	* *	56	* *
New Hope	1	* *	0	* *	2	* *
Newaygo	242	58.7 (26)	148	19.6 (21)	123	14.6 (21)
Newberry C	58	* *	44	* *	48	* *
North Camp	119	92.4 (2)	103	66.0 (1)	88	* *
Northeast	768	33.5(110)	700	10.6 (80)	476	8.4 (70)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 13
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Northern M	42	* *	27	* *	23	* *
Northwest	172	23.8(117)	126	9.5 (88)	107	13.1 (26)
Oakland Co	1,433	41.6 (90)	1,054	14.9 (46)	862	9.4 (63)
Oceana Co	166	53.6 (49)	133	22.6 (12)	100	17.0 (12)
Ogemaw	143	44.1 (83)	127	5.5(110)	98	* *
Olhse Star	247	50.6 (61)	139	30.2 (5)	94	* *
Olhse WIC	59	* *	41	* *	38	* *
Omni Care	0	* *	2	* *	8	* *
Omnicare	0	* *	1	* *	0	* *
Omnicare-M	13	* *	31	* *	27	* *
Ontonagon	37	* *	20	* *	18	* *
Oscoda	51	* *	37	* *	30	* *
Otsego Co	118	56.8 (37)	133	17.3 (31)	146	17.1 (11)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 14
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Park Medic	3	* *	12	* *	22	* *
Park Medic	0	* *	0	* *	0	* *
Park Medic	6	* *	10	* *	10	* *
Paw Paw Cl	131	45.8 (79)	93	* *	91	* *
PCOC	55	* *	48	* *	38	* *
Pinconning	39	* *	28	* *	19	* *
Port Huron	768	39.5 (99)	606	10.2 (82)	433	7.9 (76)
Portland	36	* *	29	* *	10	* *
Presque Is	65	* *	60	* *	55	* *
Profession	7	* *	10	* *	17	* *
Pullman	213	56.8 (37)	179	17.3 (31)	153	17.0 (12)
Reed City	153	62.1 (18)	118	11.9 (65)	87	* *
Region II	525	43.8 (84)	466	12.9 (57)	287	10.8 (49)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 15
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Riverview	19	* *	29	* *	32	* *
Roosevelt-	58	* *	27	* *	23	* *
Roscommon	129	56.6 (39)	110	7.3(102)	102	9.8 (60)
S & S Medi	1	* *	3	* *	2	* *
Saginaw Co	1,025	40.0 (95)	585	9.1 (92)	648	6.2 (92)
Saginaw-Ch	9	* *	9	* *	5	* *
Sanilac Co	264	52.7 (52)	185	12.4 (61)	161	10.6 (52)
Sault Ste	201	57.7 (33)	153	27.5 (7)	118	16.1 (18)
Sawyer Afb	66	* *	51	* *	38	* *
Schoolcraf	437	39.8 (96)	340	8.5 (96)	261	7.3 (85)
Seton Cent	5	* *	23	* *	30	* *
Shaftsburg	33	* *	21	* *	26	* *
Shelby M	85	* *	68	* *	78	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 16
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Sheldon Co	519	49.5 (69)	451	12.9 (57)	392	5.4 (95)
Shiawassee	335	56.4 (40)	207	11.1 (73)	223	10.3 (56)
Southfield	875	50.7 (60)	733	17.6 (29)	538	13.0 (28)
Sparta Pub	34	* *	62	* *	45	* *
St Charles	65	* *	41	* *	27	* *
St Ignace	53	* *	46	* *	44	* *
St Johns W	123	49.6 (66)	110	14.5 (50)	72	* *
St Louis C	58	* *	50	* *	25	* *
St.Clair S	512	42.6 (88)	400	11.5 (68)	301	8.0 (74)
Stanton Cl	224	49.6 (66)	180	13.9 (54)	122	13.1 (26)
Stockbridg	6	* *	13	* *	10	* *
Sumpter He	352	47.4 (74)	281	11.7 (66)	196	12.2 (40)
Taylor	873	38.4(105)	467	10.1 (84)	403	8.7 (67)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 17
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Teen Healt	1	* *	2	* *	1	* *
Temperance	5	* *	6	* *	10	* *
Thompsonv	8	* *	7	* *	11	* *
Three Oaks	45	* *	44	* *	29	* *
Three Rive	297	46.1 (77)	197	10.2 (82)	143	7.0 (87)
Traverse C	451	73.8 (5)	383	29.5 (6)	302	23.2 (3)
Tri-Cnty H	178	52.8 (51)	124	17.7 (28)	121	10.7 (51)
Tuscola	399	56.9 (36)	307	9.4 (89)	236	8.9 (65)
U Of M-Ped	6	* *	2	* *	3	* *
Vcms River	17	* *	32	* *	20	* *
Vicksburg	28	* *	17	* *	20	* *
Walled Lak	383	52.0 (54)	261	17.6 (29)	170	11.8 (44)
Warren Cli	770	39.1(100)	610	7.7(101)	444	5.9 (94)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 18
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Washtenaw	29	* *	24	* *	16	* *
Wayne-Dear	786	74.0 (4)	483	49.3 (2)	478	40.4 (1)
Webbervill	13	* *	10	* *	7	* *
West Genes	109	50.5 (62)	148	10.1 (84)	128	9.4 (63)
Western Wa	590	38.5(103)	362	10.8 (79)	311	13.2 (24)
Westland C	770	40.8 (92)	627	11.5 (68)	425	7.5 (80)
Westside C	440	61.4 (21)	402	21.6 (16)	351	16.8 (15)
Westwood	23	* *	12	* *	20	* *
Wexford	295	64.4 (16)	197	12.2 (64)	147	12.9 (30)
Wheeler WI	11	* *	9	* *	0	* *
Whitmore L	15	* *	11	* *	9	* *
Willow	1	* *	7	* *	0	* *
Woodside	57	* *	39	* *	41	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 19
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Wyoming Cl	548	59.3 (24)	496	14.7 (48)	367	12.8 (31)
Yale Clini	47	* *	34	* *	21	* *
Ypsilanti	834	54.2 (45)	612	24.5 (9)	466	18.0 (8)
Michigan	53,953	48.0	41,545	14.7	34,040	11.8
Nation (Prior Year)	969,228	52.5	527,765	20.8	555,990	12.3

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Agape WIC	1,229	11.3 (150)	7.2 (37)	4,792	9.6(20	4.5(13	13.6(16	1,899	11.6(22)	9.1 (30)		2,981	20.2 (179)
Albion Cli	521	8.6 (91)	8.8 (81)	2,102	8.8(18	5.2(15	10.9(69)	912	16.8(11	12.4(94)		1,382	12.0 (108)
Alcona	70	* *	* *	601	5.3(41)	12.5(23	6.8 (1)	295	7.1 (3)	4.4 (1)		406	9.9 (82)
Algonac Cl	337	8.6 (91)	8.0 (58)	1,354	5.7(61)	4.8(13	10.9(69)	627	17.5(14	14.5(15		942	5.0 (24)
Allegan	252	8.7 (97)	11.1 (142)	1,085	6.3(85)	6.6(19	12.6(12	456	13.4(36)	11.4(68)		736	11.3 (102)
Alma WIC	166	4.2 (6)	8.4 (69)	778	10.2(21	1.9(12)	17.7(21	343	22.7(20	21.3(19		509	10.6 (96)
Alpena Cou	527	6.5 (34)	12.3 (157)	2,408	5.6(54)	2.0(14)	13.1(14	1,132	17.1(12	12.1 (79)		1,727	9.5 (80)
Antrim - B	130	13.8 (165)	4.6 (6)	726	6.9(12	5.4(15	15.0(18	353	15.0(62)	15.9(18		504	4.6 (21)
Arab/Chald	209	7.7 (63)	8.1 (61)	1,963	5.7(61)	4.0(10	15.9(20	987	16.7(10	17.3(19		1,336	19.5 (176)
Arenac Cou	314	9.9 (127)	10.8 (134)	1,261	7.9(16	4.2(11	11.1 (78)	474	12.7 (31)	11.0(60)		891	7.2 (46)
Associated	22	* *	* *	110	2.7 (2)	5.5(16	12.7(12	49	* *	* *		77	* *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)						
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Baldwin	174	10.9 (148)	5.7 (18)	551	8.5(17	3.8(99)	9.4(34)	215	13.5(38)	12.1 (79)		311	12.2 (111)
Bangor	885	8.4 (84)	7.9 (54)	5,244	7.7(15	4.4(12	16.1(20)	2,502	17.3(13	16.2(18		3,678	13.3 (128)
Baraga Cl	111	8.1 (79)	16.2 (174)	572	4.7(25)	4.4(12	11.0(74)	295	8.5 (7)	8.8(25)		372	1.6 (3)
Barry-Eato	1,020	8.0 (77)	9.5 (101)	4,175	9.7(20	3.3(78)	12.9(13)	1,725	21.5(20	13.7(12		2,755	5.7 (29)
Barry-Eato	757	8.9 (102)	7.7 (49)	3,343	6.3(85)	5.7(17	11.8(96)	1,498	18.1(15	12.6(10		2,348	10.5 (90)
Battle Cre	2,491	8.4 (84)	8.9 (87)	8,985	8.9(19	3.8(99)	13.0(14)	3,623	16.6(10	12.8(10		5,737	18.4 (172)
Baxter Com	61	* *	* *	374	4.3(19)	2.4(33)	12.0(10)	208	14.9(60)	15.4(17		308	21.8 (188)
Bay Cl	1,894	7.7 (63)	11.1 (142)	8,007	8.2(17	2.5(40)	13.0(14)	3,628	15.6(74)	14.4(15		5,388	12.1 (110)
Bayside WI	24	* *	* *	222	8.6(18	1.4 (2)	15.8(19)	140	16.4(10	12.1 (79)		189	18.5 (173)
Beaver Isl	3	* *	* *	21	* *	* *	* *	11	* *	* *		13	* *
Belding Cl	245	9.4 (118)	8.6 (74)	962	6.9(12	3.7(98)	11.7(92)	437	20.6(19	14.2(14		668	7.9 (52)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 3
Run Date: 3 / 29 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)						
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Bell Memor	233	7.7 (63)	12.4 (159)	1,043	7.5(14	2.2(23)	16.3(20	492	18.7(16	17.9(19		674	7.4 (47)
Bendle/Bur	108	7.4 (54)	5.6 (16)	587	6.1 (74)	7.8(21	13.6(16	285	17.5(14	11.9 (75)		431	22.0 (190)
Benton Har	1,588	10.7 (143)	6.4 (24)	7,559	6.9(12	2.8(52)	12.0(10	3,445	14.6(55)	10.7 (50)		5,654	33.3 (211)
Benzie Cli	268	12.7 (161)	10.1 (114)	1,146	9.0(19	5.5(16	14.7(18	513	17.0(12	16.2(18		761	9.2 (71)
Berrien -	838	7.3 (49)	10.7 (132)	3,529	6.1 (74)	1.7 (6)	12.8(13	1,504	15.8(81)	11.1 (62)		2,464	13.6 (136)
Bldhd Migr	9	* *	* *	61	* *	* *	* *	30	* *	* *		45	* *
Branch-Hil	812	8.1 (79)	10.1 (114)	3,218	7.3(13	4.4(12	13.1(14	1,348	16.7(10	14.0(13		2,045	8.8 (67)
Branch-Hil	503	7.6 (59)	10.3 (123)	2,856	6.0(72)	4.4(12	10.9(69)	1,304	18.3(15	14.5(15		2,152	4.8 (23)
Branch-Hil	698	8.0 (77)	7.3 (40)	3,105	8.0(16	4.4(12	20.1(22	1,313	20.1(18	28.1(21		2,141	14.6 (148)
Browning-C	64	* *	* *	231	10.4(21	7.4(21	8.7(20)	86	* *	* *		159	6.9 (43)
Bryant Com	41	* *	* *	243	5.3(41)	9.1(22	10.3(47)	121	9.1 (8)	12.4(94)		167	15.0 (154)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)						
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Burton	1,727	9.3 (116)	8.5 (73)	6,413	4.9(30)	5.6(17)	10.8(64)	2,413	16.9(11)	13.5(12)		3,756	20.8 (181)
Calumet WI	220	5.0 (11)	10.9 (139)	1,156	8.0(16)	4.1(11)	13.9(16)	576	21.9(20)	14.4(15)		720	2.4 (5)
Capac Clin	96	* *	* *	396	7.1(13)	5.8(17)	15.2(19)	194	15.5(70)	20.1(19)		276	5.4 (27)
Caro WIC S	9	* *	* *	70	* *	* *	* *	38	* *	* *		60	* *
Cedar Spri	514	5.8 (25)	11.3 (147)	2,107	5.3(41)	3.2(69)	8.6(16)	918	17.1(12)	10.7(50)		1,426	1.5 (2)
Charlevoix	400	7.5 (55)	9.3 (95)	1,873	4.7(25)	4.2(11)	13.1(14)	873	18.7(16)	14.8(15)		1,386	5.1 (26)
Chass Heal	2,542	7.3 (49)	10.0 (110)	9,841	7.3(13)	5.9(18)	12.0(10)	4,029	11.1 (18)	11.8(74)		6,687	19.0 (174)
Cheboygan	371	6.2 (32)	10.2 (119)	2,036	4.7(25)	3.9(10)	11.6(90)	1,006	18.4(16)	15.5(17)		1,498	12.7 (122)
Cherry Str	677	5.8 (25)	9.0 (88)	2,684	3.5 (8)	5.3(15)	11.8(96)	1,116	15.1 (65)	16.7(18)		1,922	10.0 (83)
Chesaning	88	* *	* *	450	7.1(13)	3.6(90)	12.7(12)	207	15.9(85)	11.1 (62)		346	2.9 (10)
Clare Coun	565	9.7 (123)	6.0 (21)	2,472	6.4(96)	4.0(10)	10.8(64)	1,127	16.8(11)	12.3(88)		1,667	12.3 (112)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 5
Run Date: 3 / 29 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Clinica Sa	839	5.7 (23)	7.2 (37)	2,723	10.5(22)	3.3(78)	19.2(21)	1,039	19.2(17)	24.5(20)		1,881	12.4 (113)
Corner Hea	537	9.3 (116)	6.9 (30)	1,524	5.0(31)	7.7(21)	11.2(81)	390	17.7(15)	10.8(54)		899	20.0 (178)
Crawford	287	5.6 (19)	10.5 (128)	1,202	5.1 (36)	2.4(33)	9.9(40)	549	14.8(57)	13.3(12)		816	9.2 (71)
Dewitt WIC	133	5.3 (15)	6.8 (28)	690	7.5(14)	3.5(83)	14.1(17)	303	12.5(30)	13.9(13)		437	10.5 (90)
Dickinson-	500	5.6 (19)	14.6 (173)	2,267	6.5(10)	2.1 (18)	12.2(11)	1,072	17.3(13)	13.0(11)		1,661	7.6 (48)
Dickinson-	214	5.6 (19)	13.6 (169)	959	5.3(41)	4.1(11)	13.1(14)	467	21.2(20)	15.6(17)		678	5.8 (30)
DMC Prim	27	* *	* *	173	8.7(18)	6.9(20)	15.6(19)	87	* *	* *		127	32.3 (210)
DMC Prim	82	* *	* *	557	6.6(10)	5.4(15)	17.4(20)	293	13.7 (42)	14.0(13)		462	30.5 (207)
DMC Prim	13	* *	* *	70	* *	* *	* *	30	* *	* *		54	* *
Dorr Outre	257	7.8 (69)	10.5 (128)	1,119	8.8(18)	4.0(10)	14.7(18)	487	19.9(18)	13.1(11)		795	15.2 (156)
Douglas He	2,048	9.9 (127)	7.5 (42)	8,611	7.2(13)	5.9(18)	12.2(11)	3,702	11.8(24)	8.5(18)		5,677	13.3 (128)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 6
Run Date: 3 / 29 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Dowagiac	966	10.6 (142)	8.4 (69)	4,080	7.4(14	6.1(19	11.3(83)	1,829	13.0(33)	10.9(58)		2,833	12.4 (113)
Durand WIC	266	10.5 (138)	7.9 (54)	1,086	7.6(14	5.6(17	10.1(44)	512	16.0(92)	12.3(88)		792	4.0 (19)
East Area	732	9.7 (123)	4.6 (6)	3,524	7.8(15	6.0(18	13.6(16	1,638	15.1(65)	10.1(41)		1,145	22.9 (196)
Eastside H	1,370	15.1 (171)	3.6 (2)	6,076	9.1(19	8.6(22	12.2(11	2,647	7.1 (3)	5.4 (5)		4,383	22.8 (195)
Eau Claire	427	5.2 (13)	10.8 (134)	1,813	14.0(22	4.8(13	20.8(22	790	15.2(67)	22.3(20)		1,217	16.4 (164)
Emmet Coun	541	9.6 (121)	7.9 (54)	2,582	5.4(49)	3.4(81)	13.3(15)	1,169	16.4(10	12.8(10)		1,827	8.8 (67)
Escanaba C	768	7.7 (63)	13.0 (166)	3,438	6.2(81)	2.7(45)	10.7(60)	1,650	17.9(15	14.1(14		2,450	3.6 (16)
Family Com	574	11.5 (153)	7.0 (33)	2,521	6.6(10	6.9(20	12.4(12	1,082	13.5(38)	13.7(12		1,467	22.1 (192)
Family Hlt	628	11.6 (154)	8.3 (67)	2,475	8.0(16	7.0(20	10.8(64)	1,063	15.9(85)	11.2(65)		1,552	23.5 (198)
Family Roa	854	16.4 (175)	6.9 (30)	2,290	5.3(41)	10.0(22	10.0(42)	725	14.5(54)	11.2(65)		1,371	35.3 (212)
Farmington	116	5.2 (13)	9.5 (101)	523	6.3(85)	5.9(18	8.0 (8)	255	16.5(10	7.8(15)		358	12.0 (108)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
Fife Lake	7	* *	* *	29	* *	* *	* *	12	* *	* *	21 * *
Flint Fam	611	10.5 (138)	6.5 (26)	2,594	7.4(14)	4.2(11)	12.9(13)	1,073	16.0(92)	14.1(14)	1,733 24.8 (199)
Fruitport	21	* *	* *	76	* *	* *	* *	40	* *	* *	56 * *
Gateway	858	15.4 (172)	4.8 (9)	4,478	8.5(17)	5.4(15)	14.8(18)	2,141	16.5(10)	15.2(16)	2,791 22.6 (194)
Genesee -	2,353	10.8 (144)	7.3 (40)	11,289	5.7(61)	4.2(11)	11.0(74)	4,982	15.8(81)	11.5(70)	7,978 15.3 (157)
Gladwin Co	436	7.1 (44)	10.1 (114)	2,133	9.1(19)	1.4 (2)	13.3(15)	1,003	18.8(16)	12.3(88)	1,549 6.1 (34)
Gogebic Cl	235	6.4 (33)	7.7 (49)	1,037	8.3(17)	3.5(83)	9.2(29)	486	14.0(44)	10.9(58)	666 11.7 (106)
Golden	73	* *	* *	433	4.6(23)	6.2(19)	10.6(57)	224	8.0 (5)	7.6(12)	354 36.4 (214)
Grace Prof	252	7.5 (55)	7.5 (42)	975	3.2 (6)	10.4(22)	7.7 (5)	422	10.9(15)	5.5 (6)	675 29.6 (205)
Grace Ross	2,201	12.9 (163)	5.6 (16)	8,080	7.8(15)	4.8(13)	11.3(83)	2,901	18.5(16)	12.3(88)	5,184 14.2 (143)
Grand Have	637	8.8 (98)	9.9 (109)	2,684	7.7(15)	4.9(14)	14.5(17)	1,169	19.8(17)	16.2(18)	1,813 8.3 (61)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 8
Run Date: 3 / 29 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
Grand Trav	12	* *	* *	126	11.9(22)	6.3(19)	8.7(20)	78	* *	* *	105 14.3 (144)
Grant	283	7.8 (69)	12.0 (154)	1,242	10.4(21)	3.2(69)	15.2(19)	599	17.2(12)	14.2(14)	832 12.6 (119)
Gratiot	1,497	14.2 (167)	5.8 (19)	5,922	8.7(18)	4.6(13)	11.8(96)	2,410	14.1 (47)	8.7(23)	3,333 15.6 (159)
Greenville	309	7.8 (69)	10.0 (110)	1,199	6.7(11)	2.9(55)	12.2(11)	491	18.5(16)	10.0(38)	825 8.0 (54)
Hamtramck	413	10.2 (132)	10.7 (132)	1,988	5.0(31)	9.6(22)	9.6(35)	841	10.7(12)	8.8(25)	1,379 12.5 (117)
Hancock Cl	404	5.4 (17)	13.1 (167)	1,831	4.7(25)	4.0(10)	8.8(22)	847	9.2 (9)	5.1 (3)	1,163 3.0 (11)
Hannahvil	43	* *	* *	214	3.7(14)	0.9 (1)	19.6(22)	109	27.5(21)	20.2(19)	137 12.4 (113)
Harper Gra	2,636	14.8 (170)	5.3 (12)	9,320	10.3(21)	10.5(22)	9.6(35)	3,461	7.0 (2)	4.6 (2)	5,262 21.3 (185)
Haslett Cl	94	* *	* *	479	5.4(49)	5.2(15)	10.6(57)	205	24.9(20)	12.7(10)	337 10.4 (88)
Haslett St	16	* *	* *	60	* *	* *	* *	9	* *	* *	34 * *
Herman Kie	2,458	14.1 (166)	4.6 (6)	8,789	7.8(15)	8.8(22)	8.9(25)	3,430	6.3 (1)	5.5 (6)	5,730 19.1 (175)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Highland P	238	12.6 (160)	6.7 (27)	1,226	7.4(14	5.3(15	8.6(16)	548	13.3(35)	8.4(17)		883	13.4 (132)
Holland Cl	1,859	10.0 (130)	8.7 (77)	9,473	8.7(18	2.6(42)	14.4(17	4,163	16.5(10	14.1(14		6,676	5.6 (28)
Holly WIC	90	* *	* *	414	9.9(21	4.6(13	9.7(37)	187	17.6(14	10.7(50)		273	8.1 (56)
Holton	67	* *	* *	351	5.7(61)	4.3(12	8.3(12)	184	17.4(13	9.2(32)		270	13.7 (139)
Howard Cit	144	5.6 (19)	10.4 (125)	630	9.4(20	1.4 (2)	14.6(18	285	21.1(19	15.1(16		445	9.0 (70)
Hudsonv	411	7.8 (69)	12.4 (159)	2,263	9.8(20	5.6(17	10.7(60)	1,049	14.6(55)	7.5(11)		1,645	5.8 (30)
Hurley Cli	477	11.3 (150)	7.5 (42)	1,935	5.8(68)	7.8(21	8.5(15)	697	10.6(11)	8.6(21)		1,359	21.1 (184)
Huron Coun	618	6.0 (29)	12.9 (165)	2,932	5.7(61)	3.9(10	10.9(69)	1,400	15.0(62)	10.5(44)		2,112	14.3 (144)
Icchn-Spar	212	8.5 (89)	9.4 (97)	1,062	12.1(22	4.0(10	20.5(22	520	17.3(13	19.4(19		855	10.3 (86)
Ingham Cou	2,600	11.2 (149)	8.6 (74)	13,255	7.6(14	5.1(15	11.5(88)	5,440	17.1(12	13.1(11		9,298	12.6 (119)
Intervale	144	9.0 (107)	7.6 (47)	1,221	9.9(21	3.3(78)	23.5(22	706	23.4(20	24.1(20		1,007	45.2 (221)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)						
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Ionia Coun	794	9.1 (110)	8.7 (77)	3,169	6.5(10)	5.9(18)	10.4(50)	1,324	20.1(18)	13.5(12)		2,191	8.4 (62)
Iosco	335	6.6 (35)	11.0 (140)	2,008	6.3(85)	4.0(10)	15.0(18)	906	20.2(19)	14.9(16)		1,400	12.7 (122)
Isabella C	934	8.4 (84)	10.6 (131)	3,781	7.0(12)	3.2(69)	11.4(86)	1,561	15.6(74)	14.5(15)		2,608	8.1 (56)
Ithaca Cli	323	7.7 (63)	8.7 (77)	1,257	8.5(17)	1.8(10)	18.0(21)	513	22.8(20)	23.8(20)		745	6.4 (39)
Ithaca WIC	0	* *	* *	4	* *	* *	* *	3	* *	* *		4	* *
Jackson Co	3,266	8.6 (91)	8.6 (74)	12,658	7.7(15)	6.6(19)	11.7(92)	5,343	17.3(13)	14.0(13)		7,924	6.9 (43)
Janes St A	1,074	14.7 (169)	5.5 (15)	4,930	7.8(15)	2.4(33)	17.3(20)	2,578	15.6(74)	12.8(10)		3,663	19.7 (177)
Joy-Greenf	2,006	10.8 (144)	7.9 (54)	8,530	5.6(54)	6.6(19)	10.4(50)	3,826	14.2(49)	9.9(37)		5,523	26.0 (200)
Kalamazoo	2,280	8.9 (102)	10.0 (110)	7,978	10.3(21)	7.3(21)	11.4(86)	3,351	18.1(15)	14.2(14)		4,407	17.9 (170)
Kalamazoo	900	10.2 (132)	7.7 (49)	4,308	8.2(17)	6.7(20)	12.4(12)	2,082	15.3(68)	13.7(12)		3,134	30.0 (206)
Kalkaska	414	8.5 (89)	9.7 (106)	1,573	8.1(17)	3.1(61)	11.0(74)	652	16.9(11)	12.4(94)		1,088	13.8 (140)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Kent Count	2,873	10.8 (144)	9.4 (97)	13,399	8.2(17	2.7(45)	12.8(13	5,974	15.9(85)	11.9(75)		9,394	2.7 (7)
Kentwood	1,563	8.6 (91)	9.5 (101)	6,297	6.3(85)	3.1(61)	11.7(92)	2,591	15.6(74)	12.3(88)		4,087	2.7 (7)
Keweenaw	70	* *	* *	329	3.6(10)	2.7(45)	14.9(18	167	16.2(98)	15.0(16		231	13.4 (132)
Kingsley	33	* *	* *	229	7.4(14	2.6(42)	15.3(19	128	20.3(19	15.6(17		178	12.4 (113)
Lac Vieux	25	* *	* *	112	3.6(10)	1.8(10)	13.4(15	57	* *	* *		69	* *
Lake Co	225	8.9 (102)	10.2 (119)	1,023	7.1(13	5.1(15	10.8(64)	493	20.1(18	10.8(54)		696	14.5 (147)
Lake Odess	47	* *	* *	142	7.0(12	6.3(19	12.0(10	54	* *	* *		85	* *
Lapeer	1,323	7.3 (49)	11.2 (145)	5,166	9.1(19	3.5(83)	12.9(13	2,205	20.0(18	16.0(18		3,505	9.2 (71)
Leelanu Cl	146	5.5 (18)	14.4 (172)	741	6.1(74)	3.1(61)	16.2(20	336	14.0(44)	16.1(18		516	10.5 (90)
Leslie-Sto	55	* *	* *	293	7.2(13	7.8(21	12.6(12	123	20.3(19	18.7(19		212	8.5 (64)
Lincoln Pa	1,219	8.1 (79)	7.8 (53)	4,145	6.7(11	2.3(28)	12.0(10	1,454	18.3(15	11.9(75)		2,620	8.7 (65)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Livingston	890	7.9 (75)	12.8 (164)	3,373	5.5(53)	3.9(10)	11.9(10)	1,314	11.9(27)	8.5(18)		2,264	3.8 (17)
Lowell	97	* *	* *	490	5.1(36)	2.4(33)	11.0(74)	224	14.3(50)	12.1(79)		314	1.0 (1)
M-72 Migra	11	* *	* *	123	14.6(23)	8.1(21)	19.5(22)	61	* *	* *		103	27.2 (201)
Mancelona	201	3.5 (4)	9.5 (101)	950	5.7(61)	4.6(13)	12.4(12)	470	17.0(12)	14.7(15)		700	2.7 (7)
Manchester	5	* *	* *	54	* *	* *	* *	38	* *	* *		43	* *
Manchester	27	* *	* *	136	5.1(36)	4.4(12)	16.9(20)	63	* *	* *		91	* *
Manistee C	401	9.2 (113)	9.0 (88)	1,675	5.7(61)	3.2(69)	10.5(55)	739	17.6(14)	12.9(11)		1,169	14.8 (149)
Manistique	184	4.3 (7)	12.0 (154)	947	3.6(10)	10.0(22)	8.2(10)	493	18.1(15)	12.4(94)		677	15.4 (158)
Mansitee M	20	* *	* *	136	14.0(22)	2.2(23)	15.4(19)	76	* *	* *		105	22.9 (196)
Marion Bra	59	* *	* *	505	8.5(17)	3.0(57)	11.9(10)	237	15.6(74)	11.0(60)		391	9.2 (71)
Marquette	393	4.6 (9)	11.7 (149)	1,640	6.6(10)	2.3(28)	11.8(96)	712	17.6(14)	13.3(12)		1,052	11.1 (100)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Marshall C	262	8.4 (84)	8.8 (81)	1,039	7.9(16	3.6(90)	13.6(16	433	20.1(18	16.2(18		662	11.5 (105)
Mason Clin	86	* *	* *	360	5.3(41)	3.6(90)	12.8(13	147	14.3(50)	15.6(17		241	7.1 (45)
Mason Co	509	7.9 (75)	10.4 (125)	2,244	5.6(54)	4.4(12	9.3(32)	1,043	10.0(10)	9.6(34)		1,625	13.4 (132)
Mccree Hea	3,216	12.1 (157)	5.3 (12)	12,319	5.9(71)	4.2(11	10.7(60)	5,308	17.3(13	12.8(10		7,972	22.0 (190)
Mcnamara H	180	12.8 (162)	9.4 (97)	678	7.2(13	2.4(33)	14.6(18	315	15.9(85)	15.2(16		487	3.3 (15)
Mecosta Co	818	6.6 (35)	11.9 (152)	3,445	4.8(29)	5.4(15	7.9 (6)	1,505	14.8(57)	8.5(18)		2,362	22.2 (193)
Medical Pr	22	* *	* *	166	3.6(10)	6.0(18	12.7(12	86	* *	* *		125	31.2 (208)
Medical Pr	56	* *	* *	481	5.6(54)	6.9(20	11.6(90)	294	19.4(17	8.8(25)		383	38.6 (218)
Menominee	358	7.8 (69)	10.1 (114)	1,650	5.0(31)	2.7(45)	7.5 (4)	743	17.4(13	9.0(29)		1,117	3.0 (11)
Mercy	85	* *	* *	440	8.2(17	3.2(69)	22.3(22	200	15.5(70)	15.0(16		341	38.1 (216)
Mesick	54	* *	* *	232	9.1(19	1.7 (6)	9.1(27)	107	19.6(17	12.1(79)		160	11.3 (102)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Michigan S	130	6.9 (40)	9.2 (94)	798	4.3(19)	7.0(20)	6.9 (2)	360	13.6(41)	10.0(38)	567	9.3 (76)
Midland Cl	1,136	9.4 (118)	8.8 (81)	4,915	7.9(16)	2.3(28)	16.2(20)	2,202	15.8(81)	11.6(71)	3,589	12.8 (125)
Midwest	1,052	10.4 (135)	7.5 (42)	4,912	6.8(11)	6.4(19)	16.5(20)	2,409	17.9(15)	15.4(17)	1,583	6.3 (38)
Migrant H	221	5.0 (11)	9.0 (88)	1,122	4.5(22)	1.7 (6)	23.9(22)	584	17.5(14)	24.7(20)	794	10.3 (86)
Missaukee	205	3.9 (5)	13.7 (170)	962	5.0(31)	3.0(57)	9.9(40)	470	19.1(17)	13.2(11)	662	10.7 (98)
Moms & Bab	915	10.5 (138)	6.9 (30)	3,150	11.5(22)	5.0(14)	19.3(22)	1,183	19.1(17)	18.9(19)	1,442	9.2 (71)
Monroe Cou	1,952	9.0 (107)	7.2 (37)	7,655	7.3(13)	2.1 (18)	12.9(13)	3,225	17.4(13)	13.5(12)	4,903	17.0 (168)
Montague/	81	* *	* *	424	6.4(96)	3.1(61)	8.3(12)	208	20.2(19)	7.7(13)	310	8.7 (65)
Montcalm	21	* *	* *	110	6.4(96)	3.6(90)	16.4(20)	61	* *	* *	93	* *
Montmorenc	174	10.3 (134)	6.3 (23)	813	6.5(10)	4.9(14)	13.4(15)	407	11.8(24)	13.3(12)	590	9.3 (76)
Mt Clemens	3,478	8.8 (98)	9.0 (88)	12,679	6.3(85)	3.1(61)	13.1(14)	4,855	14.0(44)	10.3(43)	8,149	13.3 (128)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Munising C	145	6.9 (40)	11.0 (140)	655	5.2(39)	3.1(61)	13.6(16)	299	20.7(19)	20.1(19)		508	6.1 (34)
Muskegon C	3,721	10.5 (138)	8.4 (69)	15,981	6.2(81)	5.5(16)	10.1(44)	7,503	16.2(98)	12.0(78)		10,463	14.9 (152)
New Center	14	* *	* *	111	5.4(49)	4.5(13)	13.5(15)	65	* *	* *		91	* *
New Haven	299	9.0 (107)	8.7 (77)	1,068	8.3(17)	2.1 (18)	19.2(21)	381	15.5(70)	17.1(18)		665	10.5 (90)
New Hope	10	* *	* *	71	* *	* *	* *	34	* *	* *		54	* *
Newaygo	668	7.5 (55)	9.4 (97)	2,875	8.0(16)	3.5(83)	10.7(60)	1,344	12.3(29)	10.6(47)		1,834	18.2 (171)
Newberry C	161	7.5 (55)	8.1 (61)	779	6.4(96)	2.4(33)	18.2(21)	378	19.8(17)	24.9(20)		541	10.5 (90)
North Camp	315	7.0 (42)	8.3 (67)	1,289	4.1(18)	5.8(17)	7.1 (3)	512	11.7(23)	5.5 (6)		785	21.7 (186)
Northeast	2,101	14.2 (167)	5.0 (10)	9,141	9.7(20)	6.0(18)	14.2(17)	3,677	11.1 (18)	9.2(32)		6,225	15.6 (159)
Northern M	126	1.6 (2)	12.7 (163)	539	5.8(68)	2.0(14)	11.5(88)	244	19.3(17)	13.9(13)		340	7.9 (52)
Northwest	385	13.0 (164)	6.8 (28)	1,794	6.6(10)	8.2(21)	9.2(29)	835	13.5(38)	8.6(21)		820	14.8 (149)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years ⁽²⁾

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)						
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Oakland Co	4,490	9.9 (127)	8.1 (61)	18,392	6.3(85)	3.4(81)	10.5(55)	7,496	17.4(13)	13.8(13)		12,776	14.0 (142)
Oceana Co	485	6.0 (29)	9.7 (106)	2,109	7.8(15)	1.7 (6)	11.9(10)	947	18.8(16)	12.6(10)		1,480	10.7 (98)
Ogemaw	361	5.3 (15)	10.0 (110)	1,991	5.8(68)	3.6(90)	11.1 (78)	910	16.0(92)	9.8(35)		1,358	12.7 (122)
Olhsa Star	328	10.4 (135)	6.4 (24)	1,224	6.1 (74)	5.1(15)	8.6(16)	519	12.7(31)	9.1(30)		765	10.5 (90)
Olhsa WIC	114	11.4 (152)	4.4 (4)	434	9.9(21)	3.5(83)	10.4(50)	194	11.9(27)	9.8(35)		277	17.7 (169)
Omni Care	11	* *	* *	124	8.9(19)	3.2(69)	18.5(21)	71	* *	* *		96	* *
Omnicare	2	* *	* *	31	* *	* *	* *	19	* *	* *		26	* *
Omnicare-M	60	* *	* *	381	6.6(10)	5.5(16)	14.2(17)	203	15.8(81)	10.8(54)		310	27.4 (202)
Ontonagon	90	* *	* *	452	3.1 (5)	4.0(10)	8.4(14)	224	8.0 (5)	7.1 (9)		314	7.6 (48)
Oscoda	126	7.1 (44)	11.1 (142)	627	5.6(54)	4.8(13)	12.3(11)	286	16.8(11)	10.5(44)		418	11.2 (101)
Otsego Co	381	7.3 (49)	10.8 (134)	2,305	3.5 (8)	4.9(14)	10.3(47)	1,149	13.4(36)	10.6(47)		1,725	3.1 (13)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
Park Clin	2	* *	* *	12	* *	* *	* *	7	* *	* *	14 * *
Park Medic	16	* *	* *	149	3.4 (7)	4.7(13)	14.8(18)	89	* *	* *	118 40.7 (219)
Park Medic	11	* *	* *	62	* *	* *	* *	37	* *	* *	47 * *
Park Medic	36	* *	* *	209	8.6(18)	3.8(99)	25.4(23)	103	18.4(16)	28.2(21)	170 44.7 (220)
Paw Paw Cl	295	6.8 (39)	10.5 (128)	1,728	9.1(19)	5.0(14)	13.5(15)	780	15.0(62)	11.2(65)	1,206 16.8 (166)
PCOC	136	8.1 (79)	12.5 (161)	625	7.0(12)	5.6(17)	10.6(57)	272	19.9(18)	13.2(11)	350 15.1 (155)
Pinconning	101	0.0 (1)	19.8 (175)	401	6.5(10)	2.0(14)	11.7(92)	167	16.2(98)	12.6(10)	245 10.6 (96)
Port Huron	2,240	9.1 (110)	5.4 (14)	9,058	6.1(74)	8.0(21)	10.1(44)	3,985	15.4(69)	11.1(62)	6,330 8.4 (62)
Portland	89	* *	* *	305	8.2(17)	5.9(18)	14.4(17)	115	17.4(13)	26.1(20)	200 8.0 (54)
Presque Is	189	9.5 (120)	9.0 (88)	857	3.7(14)	2.2(23)	10.3(47)	390	14.4(52)	12.6(10)	629 16.9 (167)
Profession	51	* *	* *	433	2.8 (3)	5.1(15)	13.9(16)	270	14.8(57)	12.2(84)	352 38.4 (217)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 18
Run Date: 3 / 29 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Pullman	574	5.9 (27)	5.2 (11)	2,937	6.8(11	3.6(90)	13.4(15	1,437	15.6(74)	14.4(15		2,137	15.9 (163)
Ravenna	28	* *	* *	140	8.6(18	5.0(14	10.0(42)	62	* *	* *		106	29.2 (204)
Reed City	407	7.1 (44)	13.8 (171)	1,711	4.6(23)	4.9(14	10.8(64)	733	18.8(16	13.5(12		1,199	13.6 (136)
Region II	1,552	7.3 (49)	5.9 (20)	5,903	7.8(15	6.8(20	13.0(14	2,419	17.2(12	15.5(17		3,590	6.7 (41)
Riverview	52	* *	* *	246	2.4 (1)	2.8(52)	9.3(32)	121	18.2(15	7.4(10)		183	35.5 (213)
Romulus Fi	309	8.1 (79)	7.1 (36)	981	8.0(16	5.4(15	14.0(17	348	16.7(10	13.2(11		654	20.9 (182)
Roosevelt-	141	16.3 (174)	3.5 (1)	646	7.3(13	2.6(42)	15.5(19	360	16.1 (96)	10.6(47)		466	21.7 (186)
Roscommon	388	7.7 (63)	10.3 (123)	1,773	5.6(54)	4.3(12	8.8(22)	829	11.8(24)	8.8(25)		1,292	5.0 (24)
S & S Medi	20	* *	* *	166	6.6(10	3.6(90)	17.5(21	106	21.7(20	15.1(16		147	36.7 (215)
Saginaw Co	3,312	10.0 (130)	7.6 (47)	13,505	8.9(19	3.0(57)	14.3(17	6,100	17.6(14	13.7(12		9,843	8.8 (67)
Saginaw-Ch	35	* *	* *	128	12.5(22	3.1 (61)	25.8(23	40	* *	* *		78	* *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic ^(3,4)													
Clinic	Birthweight ⁽⁵⁾			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Sanilac Co	782	9.1 (110)	7.0 (33)	3,180	5.3(41)	5.6(17)	8.2(10)	1,375	17.2(12)	12.8(10)		2,131	3.8 (17)
Sault Ste	576	4.9 (10)	12.3 (157)	2,683	6.3(85)	2.1 (18)	17.5(21)	1,235	20.1(18)	15.7(18)		1,892	6.2 (36)
Sawyer Afb	164	11.6 (154)	10.4 (125)	672	6.3(85)	2.2(23)	12.9(13)	303	20.1(18)	12.2(84)		433	7.6 (48)
Schoolcraf	1,329	12.5 (159)	4.1 (3)	5,156	6.8(11)	3.8(99)	12.6(12)	2,152	13.8(43)	10.5(44)		3,623	14.8 (149)
Seton Cent	612	15.4 (172)	4.4 (4)	2,597	9.0(19)	6.3(19)	12.0(10)	969	10.7 (12)	8.7(23)		1,862	20.9 (182)
Shafsborg	79	* *	* *	308	11.4(22)	6.8(20)	14.6(18)	148	15.5(70)	23.0(20)		230	10.0 (83)
Shelby M	233	2.6 (3)	11.2 (145)	1,398	9.8(20)	2.4(33)	18.4(21)	745	21.2(20)	24.0(20)		1,130	16.7 (165)
Sheldon Co	1,635	9.8 (126)	6.1 (22)	8,214	7.1(13)	2.7(45)	15.2(19)	4,082	16.9(11)	15.1(16)		6,236	4.5 (20)
Shiawassee	1,030	7.6 (59)	8.8 (81)	3,959	8.0(16)	6.3(19)	11.9(10)	1,837	17.4(13)	14.2(14)		2,904	8.2 (59)
Southfield	2,628	10.4 (135)	8.0 (58)	11,095	6.3(85)	5.4(15)	8.0 (8)	4,362	11.3(20)	7.7(13)		7,598	13.5 (135)
Sparta Pub	290	6.6 (35)	11.4 (148)	1,327	6.6(10)	3.8(99)	9.7(37)	590	14.1 (47)	12.2(84)		929	2.2 (4)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
St Charles	158	7.0 (42)	13.3 (168)	678	10.3(21)	2.9(55)	13.3(15)	324	20.7(19)	15.4(17)		496	6.0 (33)
St Ignace	150	6.7 (38)	12.0 (154)	773	4.4(21)	3.5(83)	11.8(96)	377	19.9(18)	14.9(16)		588	4.6 (21)
St Johns W	319	7.8 (69)	8.8 (81)	1,484	5.0(31)	4.9(14)	9.1(27)	627	11.5(21)	5.1 (3)		939	12.5 (117)
St Louis C	168	8.9 (102)	11.9 (152)	631	11.6(22)	2.5(40)	17.7(21)	262	26.0(21)	23.7(20)		363	5.8 (30)
St.Clair S	1,516	9.2 (113)	8.4 (69)	5,269	6.1 (74)	3.0(57)	12.3(11)	1,945	14.9(60)	11.4(68)		3,357	13.3 (128)
Stanton Cl	653	8.6 (91)	8.1 (61)	2,710	7.9(16)	2.2(23)	13.7(16)	1,175	17.4(13)	12.4(94)		1,891	7.7 (51)
Stockbridg	36	* *	* *	212	5.2(39)	5.7(17)	12.7(12)	101	10.9(15)	18.8(19)		147	8.2 (59)
Sumpter He	904	12.2 (158)	9.6 (105)	3,385	7.2(13)	2.7(45)	11.9(10)	1,301	16.8(11)	13.2(11)		2,180	9.4 (79)
Taylor	2,521	8.9 (102)	9.3 (95)	8,800	9.5(20)	3.9(10)	11.3(83)	3,124	16.2(98)	13.8(13)		5,675	14.9 (152)
Teen Healt	12	* *	* *	86	* *	* *	* *	44	* *	* *		66	* *
Temperance	70	* *	* *	346	2.9 (4)	3.2(69)	12.1(11)	140	10.7(12)	14.3(14)		247	9.7 (81)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Comparison of Growth and Anemia Indicators by Clinic (3,4)											
Clinic	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)		≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
Thompsonv	13	* *	* *	86	* *	* *	* *	50	* *	* *	61 * *
Three Oaks	134	6.0 (29)	8.2 (66)	514	5.3(41)	1.6 (5)	11.9(10)	217	19.8(17)	13.4(12)	344 9.3 (76)
Three Rive	812	7.6 (59)	7.5 (42)	3,089	8.8(18)	1.9(12)	15.0(18)	1,258	16.9(11)	12.3(88)	2,030 8.1 (56)
Traverse C	1,214	8.6 (91)	11.8 (150)	5,078	6.8(11)	5.7(17)	12.3(11)	2,190	17.2(12)	12.4(94)	3,447 12.9 (126)
Tri-Cnty H	542	5.9 (27)	12.5 (161)	2,142	6.8(11)	5.7(17)	12.4(12)	941	18.1(15)	12.6(10)	1,450 13.1 (127)
Tuscola	1,139	7.1 (44)	9.7 (106)	4,596	6.8(11)	3.5(83)	13.3(15)	1,952	15.9(85)	13.4(12)	3,144 6.2 (36)
U Of M-Ped	15	* *	* *	71	* *	* *	* *	36	* *	* *	54 * *
Vcms River	68	* *	* *	424	3.8(16)	20.3(23)	9.0(26)	218	11.0(17)	8.3(16)	384 31.8 (209)
Vicksburg	98	* *	* *	411	9.5(20)	8.3(22)	10.9(69)	190	16.3(10)	14.7(15)	260 13.8 (140)
Walled Lak	1,073	8.8 (98)	8.8 (81)	3,710	6.2(81)	3.6(90)	10.4(50)	1,320	16.0(92)	10.0(38)	2,428 13.6 (136)
Warren Cli	2,184	9.7 (123)	8.0 (58)	8,184	6.7(11)	2.3(28)	13.7(16)	3,194	15.9(85)	12.7(10)	5,321 11.3 (102)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 22
Run Date: 3 / 29 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)												
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low		High	Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)				
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Washtenaw	75	* *	* *	413	6.1 (74)	8.2(21)	9.7(37)	181	17.7(15)	14.9(16)	257	15.6 (159)
Wayne-Dear	2,119	7.6 (59)	7.0 (33)	9,472	5.6(54)	3.2(69)	13.2(15)	4,207	15.9(85)	13.8(13)	6,665	14.3 (144)
Webbervill	40	* *	* *	178	9.0(19)	3.9(10)	7.9 (6)	79	* *	* *	119	10.1 (85)
West Genes	384	9.6 (121)	10.2 (119)	2,640	6.2(81)	5.8(17)	11.1 (78)	1,341	16.8(11)	12.2(84)	1,975	12.6 (119)
Western Wa	1,264	8.8 (98)	9.1 (93)	4,563	6.6(10)	2.7(45)	14.6(18)	1,885	18.6(16)	14.6(15)	3,018	29.1 (203)
Westland C	2,311	10.8 (144)	8.1 (61)	7,472	6.0(72)	2.8(52)	11.2(81)	2,455	17.6(14)	14.3(14)	4,571	6.7 (41)
Westside C	1,176	5.7 (23)	10.2 (119)	6,110	6.3(85)	2.0(14)	12.9(13)	2,891	19.8(17)	15.3(17)	4,532	2.4 (5)
Westwood	62	* *	* *	304	12.2(22)	4.9(14)	14.5(17)	125	14.4(52)	15.2(16)	178	20.2 (179)
Wexford	765	7.2 (48)	11.8 (150)	2,942	5.4(49)	3.2(69)	8.8(22)	1,234	15.6(74)	10.7(50)	1,935	11.8 (107)
Wheeler WI	34	* *	* *	129	7.0(12)	2.3(28)	17.8(21)	51	* *	* *	77	* *
Whitmore L	43	* *	* *	175	10.9(22)	6.9(20)	8.6(16)	72	* *	* *	101	15.8 (162)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6A
2001 - 2003 Pediatric Nutrition Surveillance (1)
Michigan
Comparison of Growth and Anemia Indicators by Clinic
Children Aged < 5 Years (2)

Page: 23
Run Date: 3 / 29 / 04

Comparison of Growth and Anemia Indicators by Clinic (3,4)													
Clinic	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Willow	25	* *	* *	105	3.8(16)	7.6(21	13.3(15	28	* *	* *		68	* *
Woodside	158	4.4 (8)	10.8(134)	755	7.7(15	2.1(18)	14.4(17	372	21.8(20	10.8(54)		531	10.4 (88)
Wyoming Cl	1,367	8.4 (84)	10.8(134)	5,557	7.1(13	3.1(61)	12.5(12	2,363	13.0(33)	11.6(71)		3,665	3.2 (14)
Yale Clini	119	11.8(156)	10.1(114)	488	6.4(96)	5.5(16	9.2(29)	216	18.5(16	10.2(42)		341	6.5 (40)
Ypsilanti	2,535	9.2(113)	7.7 (49)	10,041	6.4(96)	6.5(19	10.4(50)	4,205	16.1(96)	11.7(73)		6,539	21.8(188)
Michigan	155,916	9.5	8.3	654,410	7.2	4.5	12.2	281,480	15.8	12.5		438,342	13.4

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 1
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Agape WIC	1,499	44.5(138)	1,240	10.7(118)	852	10.9 (68)
Albion Cli	589	36.0(169)	495	7.3(169)	406	5.4(158)
Alcona	159	49.7(112)	136	16.9 (43)	99	* *
Algonac Cl	380	44.2(139)	294	12.2 (94)	241	7.5(127)
Allegan	309	54.7 (75)	266	14.7 (67)	213	8.9(102)
Alma WIC	202	44.6(137)	189	11.6(108)	146	8.9(102)
Alpena Cou	574	59.4 (40)	515	18.1 (36)	481	14.1 (33)
Antrim - B	181	74.6 (8)	159	23.9 (15)	131	16.0 (23)
Arab/Chald	472	56.1 (62)	392	21.7 (22)	337	20.8 (11)
Arenac Cou	351	52.4 (90)	337	15.7 (55)	289	11.8 (55)
Associated	29	* *	31	* *	24	* *
Baldwin	257	51.0(101)	147	17.7 (38)	86	* *
Bangor	1,330	51.7 (97)	1,167	12.8 (87)	1,005	12.9 (48)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 2
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Baraga Cl	127	53.5 (81)	117	16.2 (48)	99	* *
Barry-Eato	1,247	56.8 (55)	1,072	12.4 (91)	792	11.6 (56)
Barry-Eato	924	53.4 (82)	805	11.2(113)	660	8.2(116)
Battle Cre	2,804	41.4(151)	2,283	8.8(152)	1,672	6.5(147)
Baxter Com	79	* *	76	* *	65	* *
Bay Cl	2,135	50.5(105)	1,796	9.9(134)	1,496	6.6(146)
Bayside WI	36	* *	37	* *	44	* *
Beaver Isl	5	* *	6	* *	3	* *
Belding Cl	261	38.7(164)	216	10.2(130)	172	7.6(124)
Bell Memor	269	54.3 (77)	227	17.2 (41)	189	11.6 (56)
Bendle/Bur	134	34.3(172)	116	6.0(178)	97	* *
Benton Har	1,947	41.7(150)	2,077	8.8(152)	1,629	6.8(138)
Benzie Cli	304	70.7 (11)	265	26.0 (12)	201	17.4 (17)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 3
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Berrien -	974	48.2(119)	873	14.0 (74)	710	13.1 (45)
Bldhd Migr	14	* *	12	* *	13	* *
Booth Clin	0	* *	0	* *	0	* *
Branch-Hil	955	46.1(131)	741	12.4 (91)	576	10.4 (77)
Branch-Hil	700	47.3(124)	719	9.2(149)	575	8.7(106)
Branch-Hil	913	55.8 (65)	722	18.7 (34)	597	14.1 (33)
Browning-C	72	* *	67	* *	49	* *
Bryant Com	71	* *	64	* *	51	* *
Burton	2,754	40.9(152)	1,793	9.4(146)	1,174	7.4(131)
Calumet WI	263	58.6 (43)	237	22.8 (17)	192	17.7 (16)
Capac Clin	106	43.4(141)	102	10.8(117)	61	* *
Caro WIC S	13	* *	16	* *	12	* *
Cedar Spri	658	59.6 (38)	552	13.8 (77)	452	8.6(108)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 4
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Charlevoix	488	62.5 (25)	410	16.3 (47)	339	12.1 (54)
Chass Heal	3,031	54.0 (79)	1,958	15.7 (55)	2,091	14.9 (26)
Cheboygan	496	61.1 (29)	478	16.5 (46)	384	13.0 (47)
Cherry Str	775	65.8 (18)	654	19.4 (33)	591	13.4 (43)
Chesaning	90	* *	85	* *	80	* *
Clare Coun	665	40.8(153)	569	6.3(176)	484	6.8(138)
Clinica Sa	870	96.1 (1)	721	16.2 (48)	544	13.2 (44)
Corner Hea	556	50.0(109)	466	7.7(165)	367	6.0(153)
Crawford	315	60.3 (34)	273	11.7(105)	222	9.5 (90)
Dewitt WIC	198	57.1 (53)	173	14.5 (71)	124	13.7 (39)
Dickinson-	575	55.7 (66)	516	15.1 (63)	430	9.8 (83)
Dickinson-	240	57.1 (53)	197	13.2 (83)	156	12.2 (53)
DMC Prim	48	* *	51	* *	43	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 5
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
DMC Prim	128	28.9(180)	140	6.4(175)	111	6.3(151)
DMC Prim	17	* *	17	* *	16	* *
Dorr Outre	313	58.5 (44)	288	11.5(109)	225	11.6 (56)
Douglas He	2,341	44.7(136)	2,031	20.2 (25)	1,630	16.4 (21)
Dowagiac	1,128	53.2 (85)	918	12.1 (97)	742	9.7 (85)
Durand WIC	287	53.3 (84)	240	14.6 (68)	214	10.7 (73)
East Area	970	23.0(187)	755	6.5(174)	632	5.5(157)
Eastside H	1,680	34.8(171)	1,525	6.0(178)	1,246	5.9(155)
Eau Claire	547	81.4 (5)	450	38.7 (4)	348	27.9 (4)
Emmet Coun	724	65.5 (19)	579	21.2 (23)	456	18.9 (12)
Escanaba C	870	48.0(121)	802	12.0(100)	673	9.5 (90)
Family Com	852	43.7(140)	643	13.5 (81)	457	11.4 (61)
Family Hlt	763	50.9(102)	590	11.7(105)	432	6.7(141)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 6
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Family Roa	901	31.3(176)	649	12.0(100)	441	10.9 (68)
Farmington	180	60.0 (36)	114	32.5 (6)	83	* *
Fife Lake	4	* *	7	* *	6	* *
Flint Fam	919	37.0(166)	749	7.5(167)	479	5.0(163)
Fruitport	9	* *	8	* *	8	* *
Gateway	1,203	23.6(186)	938	5.7(181)	772	5.3(159)
Genesee -	2,143	37.6(165)	2,279	7.2(171)	1,905	5.1(161)
Gladwin Co	517	55.5 (67)	474	10.3(127)	399	7.0(134)
Gogebic Cl	269	56.5 (57)	214	10.3(127)	154	13.6 (40)
Golden	95	* *	107	3.7(185)	85	* *
Grace Prof	290	36.6(167)	274	9.5(141)	213	8.5(111)
Grace Ross	2,740	40.3(157)	2,323	9.6(138)	1,578	7.5(127)
Grand Have	761	54.8 (74)	680	15.1 (63)	537	11.0 (67)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 7
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Grand Trav	8	* *	13	* *	18	* *
Grant	344	57.6 (48)	270	15.9 (53)	244	12.3 (52)
Gratiot	1,821	28.2(182)	1,297	6.2(177)	1,202	4.5(164)
Greenville	360	47.8(122)	280	10.4(124)	219	7.8(121)
Hamtramck	410	50.0(109)	462	16.2 (48)	383	14.1 (33)
Hancock Cl	469	70.4 (12)	386	28.0 (10)	320	20.9 (10)
Hannahvil	53	* *	51	* *	44	* *
Harper Gra	3,317	25.1(185)	2,057	9.5(141)	1,874	8.0(118)
Haslett Cl	152	68.4 (14)	139	23.7 (16)	94	* *
Haslett St	25	* *	23	* *	16	* *
Herman Kie	3,121	33.2(173)	2,273	8.6(156)	1,581	6.7(141)
Highland P	239	39.3(162)	245	10.6(120)	218	9.6 (88)
Holland Cl	2,675	61.5 (27)	2,345	19.8 (28)	1,943	14.1 (33)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 8
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Holly WIC	145	42.1(145)	103	15.5 (58)	79	* *
Holton	30	* *	51	* *	53	* *
Howard Cit	171	57.3 (51)	149	19.5 (31)	106	15.1 (25)
Hudsonv	652	61.0 (30)	566	19.8 (28)	470	14.3 (30)
Hurley Cli	461	30.4(178)	491	5.5(182)	412	3.6(166)
Huron Coun	689	44.8(134)	644	7.6(166)	497	6.0(153)
Icchn-Spar	292	84.6 (4)	255	33.3 (5)	221	23.5 (8)
Ingham Cou	3,844	60.3 (34)	3,595	15.3 (60)	2,734	13.1 (45)
Intervale	238	27.3(183)	260	10.4(124)	223	11.2 (63)
Ionia Coun	876	46.2(130)	765	14.0 (74)	597	9.5 (90)
Iosco	497	40.8(153)	439	11.4(112)	366	10.4 (77)
Isabella C	1,073	62.0 (26)	952	11.9(102)	759	10.4 (77)
Ithaca Cli	403	48.9(116)	327	11.0(116)	203	8.4(112)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 9
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Ithaca WIC	0	* *	1	* *	0	* *
Jackson Co	3,616	52.3 (94)	3,199	10.2(130)	2,464	7.5(127)
Janes St A	1,144	33.1(174)	955	8.4(157)	844	5.3(159)
Joy-Greenf	2,313	52.4 (90)	1,977	26.1 (11)	1,652	25.1 (7)
Kalamazoo	2,480	42.0(148)	2,005	15.2 (62)	1,365	11.5 (60)
Kalamazoo	1,060	46.1(131)	883	8.0(161)	763	6.4(149)
Kalkaska	461	70.1 (13)	369	17.1 (42)	306	14.1 (33)
Kent Count	3,643	56.8 (55)	3,400	14.0 (74)	2,647	10.0 (81)
Kentwood	1,860	56.3 (59)	1,631	14.5 (71)	1,204	11.2 (63)
Keweenaw	76	* *	60	* *	57	* *
Kingsley	45	* *	50	* *	45	* *
Lac Vieux	25	* *	20	* *	19	* *
Lake Co	258	47.7(123)	204	8.8(152)	187	8.0(118)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 10
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Lake Odess	72	* *	46	* *	39	* *
Lapeer	1,515	53.2 (85)	1,239	13.6 (79)	930	9.0(100)
Leelanau Cl	183	80.9 (6)	169	30.8 (8)	150	26.0 (6)
Leslie-Sto	81	* *	84	* *	61	* *
Lincoln Pa	1,429	39.6(158)	1,122	9.5(141)	824	8.6(108)
Livingston	1,055	66.4 (16)	847	15.7 (55)	642	10.9 (68)
Lowell	121	62.8 (23)	116	18.1 (36)	79	* *
M-72 Migra	28	* *	31	* *	20	* *
Mancelona	227	59.0 (42)	207	15.9 (53)	198	11.6 (56)
Manchester	9	* *	8	* *	7	* *
Manchester	39	* *	39	* *	27	* *
Manistee C	443	57.3 (51)	376	14.9 (66)	321	11.2 (63)
Manistique	204	47.1(125)	194	12.9 (85)	179	8.9(102)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 11
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Mansitee M	34	* *	27	* *	29	* *
Marion Bra	125	49.6(113)	113	8.0(161)	99	* *
Marquette	437	60.9 (33)	373	20.1 (26)	306	18.3 (14)
Marshall C	287	50.9(102)	257	12.5 (88)	183	9.8 (83)
Mason Clin	106	66.0 (17)	97	* *	65	* *
Mason Co	564	58.0 (45)	499	14.6 (68)	480	9.2 (95)
Mccree Hea	4,017	29.7(179)	2,851	5.8(180)	2,135	3.8(165)
Mcnamara H	195	53.8 (80)	146	8.9(151)	106	7.5(127)
Mecosta Co	938	54.1 (78)	779	13.5 (81)	659	10.8 (71)
Medical Pr	35	* *	41	* *	33	* *
Medical Pr	56	* *	69	* *	73	* *
Menominee	456	43.2(142)	385	7.5(167)	338	3.0(167)
Mercy	116	18.1(188)	132	3.8(184)	108	0.9(169)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 12
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Mesick	47	* *	55	* *	52	* *
Michigan S	198	89.4 (3)	186	53.2 (2)	153	42.5 (1)
Midland Cl	1,290	65.0 (20)	1,160	16.0 (52)	949	12.4 (51)
Midwest	1,252	36.3(168)	1,025	10.6(120)	907	9.7 (85)
Migrant H	255	76.5 (7)	205	31.7 (7)	179	26.3 (5)
Missaukee	229	64.6 (21)	201	9.5(141)	179	7.8(121)
Moms & Bab	1,175	41.9(149)	853	16.1 (51)	683	16.4 (21)
Monroe Cou	2,265	45.7(133)	1,956	9.8(135)	1,450	8.2(116)
Montague/	42	* *	57	* *	60	* *
Montcalm	29	* *	21	* *	19	* *
Montmorenc	186	55.9 (63)	165	11.5(109)	152	9.2 (95)
Mt Clemens	4,119	49.2(115)	3,418	12.1 (97)	2,398	8.0(118)
Munising C	170	48.2(119)	150	16.7 (44)	132	14.4 (29)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 13
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Muskegon C	4,373	42.1(145)	3,388	9.6(138)	3,261	6.5(147)
New Center	24	* *	24	* *	17	* *
New Haven	354	53.4 (82)	308	20.5 (24)	211	15.2 (24)
New Hope	12	* *	13	* *	11	* *
Newaygo	792	61.0 (30)	589	17.3 (40)	514	12.6 (49)
Newberry C	178	52.2 (95)	165	12.1 (97)	152	12.5 (50)
North Camp	423	90.8 (2)	329	55.6 (1)	259	32.8 (3)
Northeast	2,732	31.8(175)	2,503	10.0(133)	1,885	7.7(123)
Northern M	141	66.7 (15)	115	25.2 (14)	96	* *
Northwest	551	30.5(177)	415	8.4(157)	319	6.3(151)
Oakland Co	5,224	42.5(143)	4,500	12.3 (93)	3,412	8.6(108)
Oceana Co	541	54.7 (75)	483	17.4 (39)	401	10.7 (73)
Ogemaw	528	50.8(104)	468	7.9(163)	376	7.4(131)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 14
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Olhsa Star	502	52.4 (90)	312	22.1 (21)	195	14.9 (26)
Olhsa WIC	169	34.9(170)	115	12.2 (94)	76	* *
Omni Care	18	* *	27	* *	28	* *
Omnicare	5	* *	5	* *	4	* *
Omnicare-M	91	* *	105	4.8(183)	71	* *
Ontonagon	105	55.2 (70)	91	* *	84	* *
Oscoda	159	61.0 (30)	135	11.9(102)	108	11.1 (66)
Otsego Co	536	59.7 (37)	519	18.7 (34)	494	16.8 (19)
Park Clin	3	* *	3	* *	3	* *
Park Medic	37	* *	36	* *	38	* *
Park Medic	5	* *	8	* *	7	* *
Park Medic	46	* *	48	* *	41	* *
Paw Paw Cl	450	50.4(108)	406	9.1(150)	332	8.7(106)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 15
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
PCOC	182	52.7 (88)	151	22.5 (19)	116	18.1 (15)
Pinconning	128	56.3 (59)	107	15.0 (65)	70	* *
Port Huron	2,478	39.6(158)	2,121	10.2(130)	1,701	6.9(137)
Portland	108	57.4 (50)	79	* *	47	* *
Presque Is	221	57.5 (49)	202	22.3 (20)	181	14.9 (26)
Profession	64	* *	69	* *	67	* *
Pullman	715	55.9 (63)	621	16.7 (44)	539	13.9 (38)
Ravenna	22	* *	25	* *	26	* *
Reed City	470	55.5 (67)	410	11.2(113)	323	9.6 (88)
Region II	1,818	46.3(128)	1,562	9.5(141)	1,065	8.3(115)
Riverview	81	* *	78	* *	57	* *
Romulus Fi	265	46.8(127)	252	8.7(155)	143	8.4(112)
Roosevelt-	163	25.2(184)	130	3.1(186)	113	2.7(168)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 16
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Roscommon	433	52.2 (95)	406	9.4(146)	353	7.6(124)
S & S Medi	14	* *	22	* *	19	* *
Saginaw Co	3,588	42.1(145)	2,810	7.9(163)	2,532	5.1(161)
Saginaw-Ch	48	* *	39	* *	29	* *
Sanilac Co	879	56.5 (57)	749	10.7(118)	567	8.8(105)
Sault Ste	665	61.2 (28)	578	25.4 (13)	466	18.9 (12)
Sawyer Afb	191	62.8 (23)	154	20.1 (26)	126	14.3 (30)
Schoolcraf	1,540	39.5(160)	1,326	9.6(138)	978	7.3(133)
Seton Cent	656	28.5(181)	698	7.2(171)	526	5.7(156)
Shaftsburg	114	59.6 (38)	81	* *	72	* *
Shelby M	310	57.7 (46)	274	22.6 (18)	278	17.3 (18)
Sheldon Co	1,876	46.9(126)	1,760	9.8(135)	1,577	6.8(138)
Shiawassee	1,152	54.9 (72)	917	9.4(146)	827	7.6(124)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 17
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Southfield	3,237	48.9(116)	2,822	13.6 (79)	2,060	10.8 (71)
Sparta Pub	271	51.7 (97)	297	12.5 (88)	251	9.2 (95)
St Charles	188	51.1(100)	133	14.3 (73)	119	6.7(141)
St Ignace	174	56.3 (59)	171	11.7(105)	155	8.4(112)
St Johns W	431	52.4 (90)	370	10.3(127)	262	9.9 (82)
St Louis C	204	52.5 (89)	174	13.8 (77)	106	14.2 (32)
St.Clair S	1,782	44.8(134)	1,459	10.5(122)	1,005	7.0(134)
Stanton Cl	725	50.5(105)	654	12.2 (94)	498	9.2 (95)
Stockbridg	53	* *	50	* *	36	* *
Sumpter He	1,152	49.4(114)	915	11.8(104)	624	9.0(100)
Taylor	3,042	39.1(163)	2,166	7.3(169)	1,586	6.7(141)
Teen Healt	6	* *	9	* *	9	* *
Temperance	73	* *	82	* *	75	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 18
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed (Rank)	Number	% Breastfed At Least 6 Months (Rank)	Number	% Breastfed At Least 12 Months (Rank)
Thompsonv	32	* *	27	* *	18	* *
Three Oaks	157	52.9 (87)	131	15.3 (60)	103	13.6 (40)
Three Rive	913	46.3(128)	753	8.4(157)	550	6.4(149)
Traverse C	1,446	73.7 (10)	1,274	28.9 (9)	1,031	21.4 (9)
Tri-Cnty H	612	49.8(111)	512	11.5(109)	392	9.7 (85)
Tuscola	1,274	54.9 (72)	1,110	10.4(124)	880	9.2 (95)
U Of M-Ped	19	* *	14	* *	16	* *
Vcms River	93	* *	103	6.8(173)	90	* *
Vicksburg	111	50.5(105)	93	* *	75	* *
Walled Lak	1,259	55.3 (69)	1,019	15.5 (58)	688	10.6 (76)
Warren Cli	2,543	40.8(153)	2,147	8.2(160)	1,592	6.7(141)
Washtenaw	94	* *	91	* *	78	* *
Wayne-Dear	2,647	74.0 (9)	1,940	40.2 (3)	1,771	33.9 (2)

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 19
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Webbervill	53	* *	40	* *	35	* *
West Genes	583	48.7(118)	561	10.5(122)	461	10.2 (80)
Western Wa	1,642	42.3(144)	1,153	12.5 (88)	833	11.3 (62)
Westland C	2,707	40.7(156)	2,226	9.8(135)	1,432	7.0(134)
Westside C	1,495	59.4 (40)	1,444	19.5 (31)	1,261	13.6 (40)
Westwood	88	* *	68	* *	66	* *
Wexford	843	63.7 (22)	664	13.0 (84)	497	10.7 (73)
Wheeler WI	38	* *	32	* *	18	* *
Whitmore L	59	* *	51	* *	33	* *
Willow	36	* *	32	* *	26	* *
Woodside	176	51.7 (97)	157	14.6 (68)	137	9.5 (90)
Wyoming Cl	1,676	57.7 (46)	1,499	12.9 (85)	1,104	9.3 (94)
Yale Clini	142	39.4(161)	117	11.1(115)	84	* *

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 7A
2001 - 2003 Pediatric Nutrition Surveillance ⁽¹⁾
Michigan

Page: 20
Run Date: 3 / 29 / 04

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ⁽²⁾
Children Aged < 5 Years ⁽³⁾

Comparison of Breastfeeding, TV Viewing, and Smoking in Household by Clinic ^(4,5)						
Clinic	Breastfeeding ⁽⁶⁾					
	Number	% Ever Breastfed	Number	% Breastfed At Least 6 Months	Number	% Breastfed At Least 12 Months
		(Rank)		(Rank)		(Rank)
Ypsilanti	3,000	55.2 (70)	2,421	19.7 (30)	1,791	16.8 (19)
Michigan	187,726	48.1	157,307	13.1	124,462	10.4

(1) Reporting period is January 1 through December 31.

(2) TV Viewing and Smoking in Household data are not reported by all contributors.

(3) Analyses based on one record per child.

(4) Excludes records with unknown data or errors.

(5) Rank compares this clinic's rate to other clinics. Rank 1 = best rate.

(6) Infants born during the reporting period included in the Ever Breastfed analysis. Infants who turned 6 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 6 Months analysis. Children who turned 12 months of age during the reporting period by/on their date of visit included in the Breastfed At Least 12 Months analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.